

Phyto-Analytical Studies and *in vitro* Wound Healing Activity of *Synedrella nodiflora* (L.) Gaertn Leaf Extract

Joshna Koodathil*, Lathika Ramadass, Sindhu Jayachandran, Jaganath Sekar

Department of Pharmaceutical Chemistry, School of Pharmacy, Sri Balaji Vidyapeeth University, Pillayarkuppam, Puducherry, INDIA.

ABSTRACT

Background: Traditionally among the Nigerian population *Synedrella nodiflora* is prominently used for treating Putrefied Wound. **Objectives:** The present work was designed to evaluate the *In vitro* Wound healing activity of *Synedrella nodiflora* leaf extracts. **Materials and Methods:** The current research work was aimed to perform the Qualitative phytochemical screening, FTIR, ICP-OES, Quantitative estimation of total phenolic and flavonoid content and *in vitro* pharmacological assay includes Antioxidant assay, Protein denaturation assay, cell line scratch assay for wound healing of EAE and EEs of *Synedrella nodiflora*. **Results:** Secondary phytochemicals like Flavonoid, Alkaloid, Tannin, Phenols, Terpenoids and Steroids were reported. The ICP-OES analysis concluded that the presence of elements like Zn, Fe, Cu, Mg and heavy metals like Ni, Pb, Cr and Cd were within their limits as per WHO. The EAE and EE of *Synedrella nodiflora* were assessed for their total phenolic and flavonoid content. The Antioxidant activity of EAE and EE showed IC_{50} value 2.24 ± 0.02 mg/mL and 3.72 ± 0.02 mg/mL compared with standard Ascorbic Acid of 42.76 ± 0.01 μ g/mL. The % Inhibition of standard Diclofenac shows IC_{50} value 27.99 ± 0.026 μ g/mL. whereas the IC_{50} values of EAE and EE were found to be 294.98 ± 0.02 μ g/mL and 138.98 ± 0.05 μ g/mL respectively. The relative cell migration of EAE and EE were 0.041 ± 0.05 mm and 0.036 ± 0.01 mm at 2mcg. **Conclusion:** These studies concluded that the *Synedrella nodiflora* leaf extracts were rich in various phytochemicals and can be used for treating wounds due to its antioxidant, anti-inflammatory and wound healing potential.

Keywords: *Synedrella nodiflora*, Phytochemical screening, Flavonoid, Phenolics, Wound Healing.

Correspondence:

Dr. Joshna Koodathil

Department of Pharmaceutical Chemistry, School of Pharmacy, Sri Balaji Vidyapeeth University, Pillayarkuppam, Puducherry, INDIA.
Email: joshnak@sbvu.ac.in

Received: 29-08-2024;

Revised: 14-11-2024;

Accepted: 22-01-2025.

INTRODUCTION

Any undesirable tissue damage resulting in the formation of wound leads to disruption of continuity towards site injury of tissues by various sets of Chemical, Physical, Immunological or Microbiological action known as biochemical reactions. Humans have localized capability of Wound healing Potential.

Plants have been considered a primary source for the treatment of various ailments since antiquity. The Secondary metabolites synthesized by plants plays a crucial role in the development of modern medicine. Globally about 80% of the population relies on classical and folk medicine for their basic needs as stated by the World Health Organization (WHO). These plant-derived medicines possess minimal or nil side effects and considered inexpensive compared to the commercial drugs available in the market.^[1]

The family *Asteraceae* is a widespread plant variety which comprises about 1600 genera and 33000 species. *Synedrella nodiflora* (L) Gaertn is a small angiospermic annual weed that grows in a distinct climatic condition in various regions such as Tropical America, Andaman, West Africa, Malaysia, India, Japan, Bangladesh, China, Spain and England. The species of *Asteraceae* is rich in phytochemicals like Alkaloids, Triterpenes, Phenols, Flavonoids, Saponin, Tannins and Polyoses. The *Synedrella nodiflora* (L) Gaertn is a small, erect annual herb. This species possesses morphological characteristics like grooved stems, dichotomous branch and swollen long internodes with hairy oppositely paired leaves up to 9 cm. The little flowers at the apex of the stem contain bunches of 2-8 efflorescence, each efflorescence with erect bracts 3-5 mm of long and floret central disc about 10-20, axillary, heterogamous. The ribbed fruits with achenes black and pappus of 2-3 bristly occurs.^[2]

The species of *Synedrella nodiflora* has a history of treating various medicinal activities like anticonvulsant, anti-inflammatory, anti-malarial, anti-diarrheal, anti-bacterial, Insecticidal, analgesic Central Nervous System (CNS) depressant and anti-psychotic activity.^[3-10] The current research work is proposed to study the Phyto analytical studies and *In vitro* Wound healing studies of *Synedrella nodiflora* leaf extracts.



DOI: 10.5530/pres.20251916

Copyright Information :

Copyright Author (s) 2025 Distributed under Creative Commons CC-BY 4.0

Publishing Partner : Manuscript Technomedia. [www.mstechnomedia.com]

MATERIALS AND METHODS

Collection of Plant Material

Synedrella nodiflora species were collected from Anumandai village Villupuram district, Tamil Nadu in the month of January during Winter season. The plant was identified and authenticated by the Auroville botanical Garden, Tamil Nadu.

Extraction

The leaves were washed by water to remove the adhered dirt and were shade dried for a period of 15 days. The dried leaves were then powdered using a mortar and pestle, passed via mesh #60 and stored in an airtight container till the study. *Synedrella nodiflora* leaf powder (30 g) was extracted by using Soxhlet extraction method for about 16 hrs with Ethyl acetate and Ethanol sequentially. The extracts were concentrated using Rotary vacuum evaporator dried extracts were kept at room temperature for further studies.^[11]

Preliminary Phytochemical Screening

The identification of Phytochemicals like terpenoids, flavonoids, alkaloids, tannins, glycosides, anthraquinones, steroids, carbohydrates and Proteins was performed using standard procedure as described in the Harborne and Biren shah with minor modifications.^[12,13]

Fourier Transform Infrared spectrophotometer (FTIR) Analysis

FTIR analysis was carried out for identification of different functional groups present in the sample Thermo Fisher (Model is 50 with In-Built ATR). A few quantities of leaf extract which was lyophilised into powdered form were encapsulated using KBr by pelletization technique. The thin film of the sample was prepared by a high pressurized system and loaded into the FTIR Spectrophotometer then the transmittance was measured between the range 400 cm⁻¹ to 4000 cm⁻¹. The samples were loaded in triplicates and bare KBr as blank. The data obtained was correlated with reference.^[14,15]

Inductively Coupled Plasma-Optical Emission Spectrometry (ICP-OES) Analysis

This ICP-OES analysis was carried out by using the instrument iCAP PRO thermo scientific make model with otegra version 2.18 (year 2021) software is used for detection and quantification of elements. About 1g of coarse leaf powder is weighed and undergoes wet digestion by using concentrated acids; add 5 mL of Conc.HNO₃, followed by 2 mL each of Con. HCl, 1 mL of 30% H₂O₂ into the vessel at 8000-10000 Kelvin to promote metallic solubilisation. To attain standard calibration curve serial dilutions from standard stock mix were prepared using distilled water (conductivity less than 2 µs/m) and the solution is directly

injected in ICP-OES. Then linear regression curve, correlation coefficient, limits of detection and quantification were determined using obtained values.^[16]

Total Phenolic content

The total Phenolic content of EAE and EE were performed by using the Folin-Ciocalteu method. About 1 mL of sample (1 mg/mL) was added to 7.5 mL of distilled water in 10 mL volumetric flask, followed by addition of 0.5 mL FC reagent and 1 mL of 35% Na₂CO₃ (Sodium Carbonate) solution into it. The sample mixture was mixed well and kept aside for 30 min and the absorbance was determined at 760 nm using a double beam UV spectrophotometer [Lasany Li 2700]. To estimate the Total Phenolic content, Gallic acid standard was used as reference. The estimated values were expressed as mg of Gallic acid equivalent (GAE/g).^[17,18]

Total Flavonoid content

Total flavonoid content of *Synedrella nodiflora* leaf extracts were measured spectrophotometrically by using Aluminium chloride method as per standard procedure. About 0.5 mL concentration of the sample extract was mixed with 7.5 mL of ethanol, 0.5 mL of 1M sodium acetate, 0.5 mL of 10% aluminium chloride and 16 mL of distilled water. The solutions were maintained at room temperature for 30 min and the absorbance was measured at 415 nm using a double beam UV spectrophotometer [Lasany Li 2700]. The results were expressed as mg/g Quercetin equivalents (QE/g).^[19,20]

Anti-oxidant activity

The antioxidant activity of the leaf extracts of *Synedrella nodiflora* were studied using free radical scavenging efficiency by 2,2-diphenyl-1-Picrylhydrazyl (DPPH) assay method. About 1 mL various concentrations of both leaf extracts (1-5 mg/mL) were added to 6 mL methanolic solution of 0.1M DPPH. Then the mixture was shaken vigorously and kept aside for 30 min at 27°C. The absorbance was measured at 517 nm using UV-Spectrophotometer [Lasany Li 2700]. Similarly, various concentrations of Ascorbic acid (10-50 µg/mL) were used as a reference standard. The percentage of Inhibition was estimated by the given formula.^[21-23]

$$\% \text{ of Free Radical inhibition} = \frac{C_A - T_A}{C_A} \times 100$$

C_A. Control absorbance;

T_A. Test sample absorbance.

In vitro Anti-inflammatory activity

In vitro Anti-inflammatory activity of *Synedrella nodiflora* leaf extracts were studied using egg albumin protein denaturation method. A successive dilution of leaf extracts ranging from 78.125 µg/mL to 2500 µg/mL was prepared and the reaction mixtures 0.4

mL of fresh egg's albumin and 5.6 mL of Phosphate buffer (adjust pH 6.4) were added.

Different concentrations of plant extract (4 mL) were mixed thoroughly with the above mixture. Various concentrations of reference standard ranges from 31.25 µg/mL to 1000 µg/mL was prepared that serves as Positive control. The denaturation process started when the reaction mixtures were kept in a water bath at 37°C±2°C initially for 15 min and then the temperature was maintained at 70°C±2°C for 3-5 min. The reaction mixture was cooled after the denaturation process for about 10-15 min. The absorbance was measured at 680 nm using a double beam UV-Spectrophotometer [Lasany Li 2700] before and after denaturation. The Percentage inhibition was measured by the given formula.^[24-27]

$$\% \text{ Inhibition} = \frac{\text{Absorbance of control} - \text{Absorbance of standard}}{\text{Absorbance of control}} \times 100$$

Cell Scratch Wound healing assay

The expansion of cell population using scratch assay was performed for the assessment of distribution and migration of 3T3-L1 cells. The cells were transferred into culture plates (6 well plates) in Dulbecco's Modified Eagle Medium (DMEM) containing 10% fetal bovine serum and 2% antibiotics like Streptomycin and Penicillin. Generation of the linear wound was done in the monolayer with the help of sterile 100 µL Plastic pipette tip, after the formation of a confluent cell monolayer. Phosphate buffer saline was used to remove cellular debris. Dimethyl Sulfoxide [DMSO] (0.01%) used as solvent control and the Positive control comprises the Platelet derived growth factor (PDGF; 0.002 µg/mL). The concentration of crude extract was taken as 1 µg/mL, 2 µg/mL. The incubation of cells was performed using 5% CO₂ for a period of 16 hr. Microscopical images (capture pro Version 2.5 for progress® microscope camera from Jenoptik laser, Optik, system GmbH) were taken after incubation of scratched cell layer for the estimation of Relative Cell Migration.^[28-32]

RESULTS

Phytochemical Screening

The preliminary phytochemical analysis of *Synedrella nodiflora* leaf extracts were represented in Table 1. The EAE and EEs revealed the presence of Alkaloids, Glycosides, Flavonoids, Terpenoids, Tannins, Phenols, Steroids and Reducing sugars.

FTIR analysis

The FTIR analysis elucidates the functional group of unknown composition of secondary metabolites that may be present in the EAE and EE which were represented in Figures 1 and 2. The spectral peak value ranges from 4000 cm⁻¹ to 1500 cm⁻¹ usually considered as functional group region and the range below 1500 cm⁻¹ to 600 cm⁻¹ may be considered as fingerprint regions for identification of traces element, alkyl or aryl bond. The presence

of compounds like ester (1736.60 cm⁻¹), phenol (1384.56 cm⁻¹), alkane methylene group (1462.94 cm⁻¹) and aldehyde (1381.81 cm⁻¹) spectral ranges may indicate the presence of aromatic compound like terpenoid class. The compound class of functional groups like delta lactone or 6 membered lactone group (1737.48 cm⁻¹), sulphonamide (1164.96 cm⁻¹) and intermolecular bonded alcohol (3402.02 cm⁻¹) may be representing the presence of cardiac glycoside. The existence of primary amine (3406.67cm⁻¹), phenols (1384.56cm⁻¹), ester group may elucidate the presence of flavonoids. The presence of conjugated alkene (1630.36cm⁻¹), carboxylic acid dimer (1710.32cm⁻¹), secondary alcohol (1113.82cm⁻¹), aldehyde (1737.48cm⁻¹), ketone (1710.32cm⁻¹) may illustrate the presence of tannins. Hence the EAE and EE may possess certain classes of secondary metabolites like terpenoid, tannin, flavonoids and glycosides as reported in Tables 2 and 3 respectively.

ICP-OES Analysis

The ICP-OES analysis result showed that the presence of some Inorganic trace elements in various concentration were Magnesium (2060.20ppm), Zinc (46.97ppm), Chromium (45.59ppm), Copper (33.08ppm), Titanium (21.91ppm), Vanadium (3.82ppm), Nickel (3.80ppm), Lead (3.55ppm) and Cadmium (0.047ppm) respectively. The Quantitative elemental detection of 11 elements was represented in Figure 3.

Total Phenolic Content

The Total Phenolic Content of EAE and EE were represented in Table 4. The Quantitative estimation of *Synedrella nodiflora* leaf extracts by using Folin-Ciocalteu Reagent is stated in terms of Gallic acid equivalent and they were expressed as mg GAE/mg. The Linear Regression Curve of Gallic acid Standard was represented in Figure 4. The EAE showed higher concentration of Phenol as compared to EE.

The data obtained were expressed as Mean±Standard Deviation (n=3).

Total Flavonoid content

The result for Total Flavonoid content of EAE and EE were represented in Table 5. The Linear Regression Curve of Quercetin Standard was represented in Figure 5. The Flavonoid quantity was estimated spectrophotometrically by the Aluminium Chloride Colorimetric method and was expressed in the terms of Quercetin Equivalent (QE/mg). The EAE exhibited higher flavonoid content as compared to EE.



Anti-oxidant activity

The extracts were tested for their free radical scavenging capacity by the DPPH assay method using ascorbic acid as a standard. The degree of scavenging activity was summarized in Table 6 and the percentage of inhibition was represented in Figures 6-8. The IC₅₀

Table 1: Result of Phytochemical analysis of *Synedrella nodiflora* leaf extracts.

Sample	Flavonoid	Alkaloid	Phenol	Glycoside	Steroid	Tannin	Terpenoid	Saponin	Reducing sugar
EAE	+++	+	+++	++	++	++	+++	-	++
EE	++	+	++	++	+	++	+++	-	++

EAE- Ethyl acetate extract; EE- Ethanolic extract. +: positive; -: negative.

**Figure 1: FTIR spectra of EAE of *Synedrella nodiflora*.****Figure 2: FTIR spectra of EE of *Synedrella nodiflora*.**

values were obtained using the Linear regression curve method and shown in Table 6.

Protein denaturation assay

The *In vitro* anti-inflammatory activity for EAE and EE of *Synedrella nodiflora* was determined by using egg albumin denaturation assay using double beam UV-spectrophotometer

[Lasany Li 2700] and the results were expressed in the Tables 7 and 8 which were Interpreted from Figures 9-11. This study proves that the percentage inhibition of protein denaturation depends upon concentration. However, the EAE shows more percentage inhibition than the standard Diclofenac, but EE shows less percentage inhibition compared with standard. The IC_{50} values of EAE, EE and standard Diclofenac were found to be

Table 2: Result of FTIR analysis on EAE of *Synedrella nodiflora*.

Sl. No.	Peak frequency cm ⁻¹	Reference Frequency Range cm ⁻¹	Functional group	Intensity of functional group	Compound class
1.	3402.02	3550-3400	N-H stretching	Medium	Primary amine
2.	2924.22	3300-2500 3000-2840	O-H stretching C-H stretching	Strong, Broad Medium	Carboxylic acid Alkane
3.	2852.30	3000-2840	N-H stretching	Strong, Broad	Amine salt
4.	1737.48	1750-1735	C=O stretching	Strong	Ester (6-membered Lactone)
5.	1710.32	1720-1705	C=O stretching	Strong	Carboxylic acid oligomer
6.	1631.20	1650-1600	C=C stretching	Medium	Conjugated alkene
7.	1554.35	1550-1500	N-O stretching	Strong	Nitro compounds
8.	1497.60	1550-1500	N-O stretching	Strong	Nitro compounds
9.	1462.94	1465	C-H bending	Medium	Alkane Methylene group
10.	1381.81	1390-1380 1390-1310 1385-1380	C-H bending O-H bending C-H bending	Medium Medium Medium	Aldehyde Phenol Alkane gem dimethyl
11.	1275	1275-1200	C-O stretching	Strong	Alkyl aryl ether
12.	1164.96	1170-1155	S=O stretching	Strong	Sulfonamide
13.	1113.82	1124-1087	C-O stretching	Strong	Secondary alcohol or Aliphatic ether
14.	1046.83	1050-1040	CO-O-CO stretching	Strong, Broad	Anhydride
15.	985.18	995-985	C=C bending	Strong	Alkene monosubstituted
16.	843.90	850-550	C-Cl stretching	Strong	Halo compound
17.	721.36	700±20	C-H bending	Strong	monosubstituted benzene derivative or Tri substituted
18.	675.99	730-665	C=C bending	Strong	Alkene Disubstituted (cis)
19.	614.57	690-515	C-Br stretching	Strong	Halo compound

138.98±0.05 µg/mL, 294.98±0.02 µg/mL and 27.99±0.026 µg/mL respectively.

Cell scratch assay method


The relative fibroblast cell migration of EAE and EE of *Synedrella nodiflora* were represented in Figure 12 and found to be 0.036 mm, 0.029 mm and 0.036 mm, 0.041 mm at lower and higher concentration respectively. The relative cell migration of standard (PDGF) was found to be 0.042mm. Those results were represented in Table 9 which showed that EAE exhibited increased Wound healing activity compared to the EE.

DISCUSSION

Evaluation of the biologically compounds present in plant extracts, present in predominantly by phytochemical screening may be helpful in the identification of therapeutically effective plant sources.^[33] Terpenoids are reported to possess different pharmacological activities including anti-inflammatory, anti-malarial, anti-cancer, anti-microbial and hypoglycemic activity.^[34] Flavonoids exhibit wide range of biological activity as anti-oxidant, anti-diabetic, anti-microbial, anti-inflammatory.^[35] Additionally, flavonoids were also recognized for its wound healing property.^[36] Alkaloids are well known for its anti-inflammatory,

Table 3: Result of FTIR analysis on EE of *Synedrella nodiflora*.

Sl. No.	Peak frequency cm^{-1}	Reference Frequency Range cm^{-1}	Functional group	Intensity of functional group	Compound class
1.	3406.67	3550-3400	N-H stretching	Medium	Primary amine
2.	2922.48	3300-2500	O-H stretching	Strong, Broad	Carboxylic acid
3.	2851.65	3000-2840	C-H stretching	Medium	Alkane
4.	1736.60	1750-1735	C=O stretching	Strong	Ester (6-membered Lactone)
5.	1630.36	1650-1600	C=C stretching	Medium	Conjugated alkene
6.	1384.56	1390-1380 1390-1310	C-H bending O-H bending	Medium Medium	Aldehyde Phenol
7.	1165.74	1165-1150	S=O stretching	Strong	Sulfonic acid
8.	1066.69	1085-1050	C-O stretching	Strong	Primary alcohol
9.	934.53	915-905	C=C bending	Strong	Alkene monosubstituted
10.	898.59	880±20	C-H stretching	Strong	1,2,4-Tri substituted
11.	825.18	810±20	C-H bending	Strong	1,2,3,4-Tetra substituted, or 1,4-Di substituted
12.	720.82	700±20	C-H bending	Strong	1,3-Disubstituted or monosubstituted benzene derivative
13.	616.65	690-515	C-Br stretching	Strong	Halo compound

**Figure 3:** ICP-OES analysis of *Synedrella nodiflora* coarse leaf Powder.

analgesic, anti-cancer activity and also for promoting wound healing.^[37] The phenolic compounds and tannins were reported to possess anti-diabetic, anti-oxidant and antibacterial activity.^[38,39] In the current study, FTIR spectroscopy is used for identifying different functional groups present in the leaf extracts

of *Synedrella nodiflora* based on the measurement of the peak values.^[40] The FTIR results of *Synedrella nodiflora* leaf extract reported the presence of groups indicating the phytochemicals like terpenoid, tannin, flavonoids and glycosides previously reported to contribute the anti-bacterial, wound healing activity.




Figure 4: Linear Regression Curve of Gallic acid Standard.




Figure 5: Linear Regression Curve of Quercetin Standard.

ICP-OES is a widely used method for element analysis due to its ability as a multiple wavelength detector for identification and quantification of elements. Among various traditional elemental analysis methods like spectroscopic, volumetric, gravimetric and colorimetric, the inductively coupled plasma methods like mass spectrometry (ICP-MS) and optical emission spectroscopy (ICP-OES) indicates higher accuracy and precision rate. Under the controlled condition ICP-OES exhibits spectral plasma excitation at different wavelengths for each distinct element. The emission of light intensity is compared with concentration of each element.^[41] Magnesium plays a vital role in initiating wound healing through different metabolic pathways and cellular mechanism. It contributes nearly 3000 enzymes for ATP, RNA and DNA formation.^[42] Iron, an important element of hemoglobin

Table 4: Total Phenolic Content of *Synedrella nodiflora* leaf extract.

Sample	Gallic acid equivalent (GAE/mg)
EAE	0.0040±0.0003
EE	0.0025±0.0001

and myoglobin. It catalyzes generated free radicals and promotes tissue injury and cellular death. According to the recommended dietary allowance, normal intake of iron for women is 18 mg/day and for men 10 mg/day. Zinc, an integral part of proteins which stabilize cellular membrane, cell respiration, wound healing, synthesis of DNA and protein and regulator homeostasis. The recommended normal level for zinc was 15 mg per day for an




Figure 6: DPPH assay of Ascorbic acid using Linear Regression method.




Figure 7: DPPH assay of EAE of *Synedrella nodiflora* using Linear Regression method.

Table 5: Total flavonoid Content of *Synedrella nodiflora* leaf extract.

Sample	Quercetin equivalent (QE/mg)
EAE	0.051±0.004
EE	0.020±0.0025

The obtained results were expressed as Mean±Standard Deviation (n=3).

adult.^[43] Copper is an essential element that plays a complex role in the mechanism of wound healing and cytokines modulators. Which tolerate the expression of collagen and fibrinogen formation The average intake of copper varies among 28 countries for adult ranging 1.6 to 3.3 mg/day.^[44] Manganese and Zinc combination used for wound healing by changing integrin expression and the average intake of manganese for adult was 2.5 to 5 mg/day.^[45] Boron is used as diet supplements because it has interaction with magnesium, Vitamin D and Calcium. It also contributes to anti-oxidant and anti-inflammatory activity.^[46] The heavy metal like lead permissible level for human adult intake

Table 6: IC₅₀ values of *Synedrella nodiflora* leaf extract using DPPH assay.

Sample	IC ₅₀ Value
Ascorbic acid	42.76±0.01 µg/mL
EAE	2.24 ±0.02 mg/mL
EE	3.72 ±0.02 mg/mL

The obtained results were performed in triplicate using Mean±Standard Deviation (n=3).

was 7 µg/kg body mass/day, according to WHO guidelines. Cadmium intake should not exceed 0.05 mg/kg according to the United States FDA from NCBI guidelines. Chromium, an essential element for carbohydrate metabolism but it comes under grade2 heavy metal. The average intake of humans differs in different regions. so, the National Research Council (1980) set the limit 200 µg/day for adults. All the elements described above were within their limit as said by WHO guidelines, FDA, NCBI guidelines and ICH guidelines. Plant phenolic compounds are considered as primary antioxidants or free scavengers and play a major role associated with the treatment of various diseases.




Figure 8: DPPH assay of EE of *Synedrella nodiflora* using Linear Regression method.




Figure 9: Protein Denaturation assay of Standard Diclofenac using Linear Regression method.

Table 7: The Percentage inhibition rate of Standard Diclofenac.

Concentration (µg/mL)	% Inhibition
31.25	25±0.01
62.5	125±0.05
125	410±0.03
250	1090±0.03
500	2100±0.04
1000	4010±0.02

Each value expresses the Mean±Standard deviation (n=3).

Table 8: The Percentage inhibition rate of *Synedrella nodiflora* leaf extract.

Concentration (µg/mL)	%Inhibition of EAE	% Inhibition of EE
78.125	20±0.001	12.5±0.001
156.5	55±0.001	25±0.003
312.5	122.6±0.005	50±0.0004
625	248.7±0.35	120±0.005
1250	505.9±0.01	210.5±0.1
2500	1000.6±0.2	456.6±0.5

Data were obtained from three independent experiments, each performed in triplicates and represented as Mean±SD.




Figure 10: Protein Denaturation assay of EAE of *Synedrella nodiflora* using Linear Regression method.




Figure 11: Protein Denaturation assay of EE of *Synedrella nodiflora* using Linear Regression method.

FC reagent method is the most employed for the quantification of phenolic compounds. The extracts containing phenolic compounds react with FC reagent in an alkaline medium forming blue coloured Phosphomolybdic/Phosphotungstic acid complex.^[47] The antioxidant activity produced by flavonoids involves inhibiting the formation of reactive oxygen and chelating the trace elements involved in the production of free radicals. The radical scavenging activity of flavonoids were reported on free radicals and oxidizing molecules and the potency depends upon the presence of OH groups. The antioxidant activity produced by flavonoids involves inhibiting the formation of reactive oxygen and chelating the trace elements involved in the production of free radicals.^[48] DPPH assay method is used to measure the electron donating capacity of natural compounds based on the free radical scavenging property. The antioxidant activity of the compound is proportional to the degree of decolourization of

the DPPH solution. A significant reduction in the absorbance of the reaction mixture indicates the higher inhibition percentage of the compound under study. In the current study the Ethyl acetate and EEs showed higher inhibition percentage may be due to the higher phenolic content.^[49] The main mechanisms involved in protein denaturation are modification of hydrogen atoms involved in electrostatic interaction, hydrophobic interaction and disulfide linkages. The production of auto antigens by denaturation is caused due to inflammations in clinical conditions like rheumatoid arthritis, cancer, rheumatoid arthritis and diabetes. The anti-inflammatory activity of a drug can be measured by its ability to inhibit denaturation of protein. Both the extracts were found to inhibit the protein denaturation by reducing the synthesis of collagen, mucopolysaccharide and reducing the number of fibroblasts.^[50] DMEM is the Dulbecco's modified eagle medium that is mainly used as growth stimulators




Figure 12: Microscopical images of *in vitro* scratch assay results of *Synedrella nodiflora* leaf extracts showing fibroblast cell migration after 24 hr of induction of Wound in the following groups a) DMSO; b) PDGF; c) EAE 1-mcg; d) EAE 2-mcg; e) EE 1-mcg; f) EE 2-mcg.

Table 9: The Relative cell migration rate of *Synedrella nodiflora* leaf extracts.

Sl. No.	Treatment Group	Concentration ($\mu\text{g/mL}$)	Relative Cell Migration (mm) at 24hr.
1.	1% DMSO		0.032 \pm 0.001
2.	PGDF	0.002	0.042 \pm 0.003
3.	EAE	1	0.036 \pm 0.0005
		2	0.041 \pm 0.0005
4.	EE	1	0.029 \pm 0.001
		2	0.036 \pm 0.001

Experimental groups were performed in triplicates and interpolated by using Mean \pm Standard Deviation ($n=3$).

in various mammalian cells. The role of DMEM in Scratch assay is regeneration of fibroblast and re-epithelization involved in the secondary phase of wound healing using Fetal bovine serum as supplement and Platelet derived growth factor (PDGF) as growth regulator during cell division. PDGF is used as positive control due to peptide matrix production, synthesis of many epithelial and connective tissues thereby increasing the efficiency of wound healing. To elucidate the effectiveness of wound healing, the wound closure time is frequently observed with a specific range of time intervals that determines cell proliferation. Some of the

studies showed that above 0.5%DMSO concentration produces toxic effects, our study has 0.01% which is negligible. For better understanding, comparative study between EAE and EE of *Synedrella nodiflora* at different concentrations with specific time intervals for cell migration and proliferation were studied with 0.02 $\mu\text{g/mL}$ of standard PDGF. The EAE shows increased relative cell migration compared to EE within 0-24 hr simultaneously. The results compared with positive control confirmed that EAE had relatively the same proliferation activity.^[51]

CONCLUSION

This is the first study reporting Wound healing potency of *Synedrella nodiflora*. The therapeutically important phytochemicals and elements like Magnesium, Iron, Zinc and Copper were reported in *Synedrella nodiflora*. All these constituents may contribute to the Wound healing activity of *Synedrella nodiflora* leaf extract. This study scientifically supported the traditional claim of *Synedrella nodiflora* for the treatment of wounds.

In the current study, the results scientifically support the traditional claim of wound healing property of *Synedrella nodiflora* plant. The results of anti-inflammatory activity of the plant extract demonstrated its potential to support wound healing. The presence of phenolic compounds, flavonoids, alkaloids, terpenes and tannins also plays a significant role in wound healing. The presence of elements like Copper, Magnesium and Zinc also plays a major role in initiating the Wound healing process. Further studies on isolating the phytochemical from *Synedrella nodiflora* will be beneficial in revealing the wound healing potential to limelight.

ACKNOWLEDGEMENT

The authors acknowledge Sri Balaji Vidyapeeth (Deemed to be University) and Synthiya Research Laboratory for their contribution in carrying out research work.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ABBREVIATIONS

EAE: Ethyl acetate extract; **EE:** Ethanol extract; **WHO:** World Health Organization; **CNS:** Central Nervous System; **FTIR:** Fourier transform infrared spectrophotometer; **ICP-OES:** Inductively Coupled Plasma-Optical Emission Spectrometry; **FC:** Folin Ciocalteu; **DPPH:** 2,2-diphenyl-1-picrylhydrazyl; **DMEM:** Dulbecco's modified Eagle Medium; **DMSO:** Dimethyl Sulfoxide; **EAE:** EAE; **EE:** EE; **FDA:** Food and drug administration; **ICH:** International council for Harmonisation; **NCBI:** National center for Biotechnology Information; **PDGF:** Platelet- derived growth factor.

SUMMARY

Development of wounds has become one of the serious concerns especially among the patients with diabetics. Phytochemical studies on *Synedrella nodiflora* leaf extract reported the presence of flavonoid, phenol, terpenoids, tannins and steroids. These phytochemicals are reported to possess Wound healing activity. The elemental analysis of *Synedrella nodiflora* leaf extract revealed the presence of the elements like Magnesium, Iron, Zinc and Copper that have a vital role in promoting Wound healing. Magnesium initiates Wound healing while Iron promotes Wound

healing by avoiding damage to the tissues and cell line death. Zinc is also an important element for promoting Wound healing by stabilizing cellular membranes and regulating Hemostasis. Copper is an essential element to produce collagen and fibrinogen there by promoting Wound healing. Terpenoids also accelerate Wound healing by promoting the production of collagen thereby promoting the Healing process. GCMS reports of *Synedrella nodiflora* leaf extracts reported the presence of high amounts of Phytol, a diterpene alcohol. Phytol reduces inflammation by inhibiting nuclear factor-Kappa B (NF-kB Pathway). It also provides better remodeling of tissue during Wound healing.

REFERENCES

- Sharma M, Sharma M, Bithel N, Sharma M. Ethnobotany, phytochemistry, pharmacology and nutritional potential of medicinal plants from *Asteraceae* Family. *J Mountain Res.* 2022;17(2):67-83. doi: 10.51220/jmr.v17i2.7.
- Y BP, J DM, S GD, Amravati S. Pharmacognostic studies and anti-microbial activity of *Synedrella nodiflora* (L.) Gaertn. 2011;2(3):317-21.
- Amoateng P, Tagoe TA, Karikari TK, Kukuia KK, Osei-Safo D, Woode E, et al. *Synedrella nodiflora* extract depresses excitatory synaptic transmission and chemically induced *in vitro* seizures in the rat hippocampus. *Front Pharmacol.* 2021;12:610025. doi: 10.3389/fphar.2021.610025, PMID 33762938.
- Abad MJ, Bermejo P, Carretero E, Martínez-Acitores C, Noguera B, Villar A. Anti-inflammatory activity of some medicinal plant extracts from Venezuela. *J Ethnopharmacol.* 1996;55(1):63-8. doi: 10.1016/s0378-8741(96)01478-x, PMID 9121169.
- Chaniad P, Techarang T, Phuwajaroanpong A, Na-ek P, Viriyavejakul P, Punsawad C. *In Vivo* Anti-malarial Activity and Toxicity Study of Extracts of *Tagetes erecta* L. and *Synedrella nodiflora* (L.) Gaertn. from the *Asteraceae* Family. *Evid Based Complement Alternat Med.* 2021; 2021:1-9.
- Zahan R, Nahar L, Haque A, Mosaddik A, Fazal A, Haque ME. Anti-diarrheal and hypoglycemic effects of *Synedrella nodiflora*. *Phytopharmacology.* 2012;2(2):257-64.
- Wijaya S, Nee TK, Jin KT, Hon LK, San LH, Wiart C. Antibacterial and antioxidant activities of *Synedrella nodiflora* (L.) Gaertn. (*Asteraceae*). *J Complement Integr Med.* 2011;8(1):1-13. doi: 10.2202/1553-3840.1499, PMID 22754934.
- Rathi M. Insecticidal activity of aerial parts of *Synedrella nodiflora* Gaertn. (Compositae) on *Spodoptera litura* (Fab.). *J Cent Eur Agric.* 2006;6(3):223-8.
- Nahar L, Zahan R, Morshed MT, Haque A, Alam Z, Mosaddik A. Antioxidant, analgesic and CNS depressant effects of *Synedrella nodiflora*. *Pharmacogn J.* 2012;4(31):29-36. doi: 10.5530/pj.2012.31.6.
- Dudev T, Lim C. Importance of Metal Hydration on the Selectivity of Mg²⁺ versus Ca²⁺ in Magnesium Ion Channels. *J Am Chem Soc.* 2013;135(45):17200-8. doi: 10.1021/ja4087769, PMID 24087890.
- Redfern J, Kinninmonth M, Burdass D, Verran J. Using Soxhlet ethanol extraction to produce and test plant material (essential oils) for their anti-microbial properties. *J Microbiol Biol Educ.* 2014;15(1):45-6. doi: 10.1128/jmbe.v15i1.656, PMID 24839520.
- Shaikh JR, Patil M. Qualitative tests for preliminary phytochemical screening: an overview. *Int J Chem Stud.* 2020;8(2):603-8. doi: 10.22271/chemi.2020.v8.i2i.8834.
- Onyeka IP, Onyegbule FA, Ibe CI, Ifebi HM, Ezugwu CO. Pharmacognostic, Acute Toxicity Profile and Comparative Leaf Characteristics of *Desmodium velutinum*, *Mallotus oppositifolius* and *Synedrella nodiflora*. *JMPPCR.* 2022;10-24. doi: 10.9734/ijmpcr/2022/v15i430163.
- Chopra H, Bibi S, Mohanta YK, Kumar Mohanta T, Kumar S, Singh I, et al. *In vitro* and *in silico* characterization of curcumin-loaded chitosan-PVA hydrogels: anti-microbial and potential wound healing activity. *Gels.* 2023;9(5):394. doi: 10.3390/gels9050394, PMID 37232986.
- Rani S, Puri R, Boora P, Qasim A, Angmo D, Mehta M, et al. FT-IR and ICP-MS Analysis of Bioactive compounds in the methanolic Leaves Extract of *Thalictrum foliolosum*. *Pharmacogn Res.* 2023;15(4):705-15.
- Masson P, Dalix T, Bussière S. Determination of major and trace elements in plant samples by inductively coupled plasma-mass spectrometry. *Commun Soil Sci Plant Anal.* 2010;41(3):231-43. doi: 10.1080/00103620903460757.
- Mustafa I, Faisal MN, Hussain G, Muzaffar H, Imran M, Ijaz MU, et al. Efficacy of *Euphorbia helioscopia* in context to a possible connection between antioxidant and antidiabetic activities: a comparative study of different extracts. *BMC Complement Med Ther.* 2021;21(1):62. doi: 10.1186/s12906-021-03237-x, PMID 33579270.
- Froehlicher T, Hennebelle T, Martin-Nizard F, Cleenewerck P, Hilbert JL, Trotin F, et al. Phenolic profiles and anti-oxidative effects of hawthorn cell suspensions, fresh fruits and medicinal dried parts. *Food Chem.* 2009;115(3):897-903. doi: 10.1016/j.foodchem.2009.01.004.
- Nurcholis W, Sya'bani Putri DN, Husnawati H, Aisyah SI, Priosoeryanto BP. Total flavonoid content and antioxidant activity of ethanol and ethyl acetate extracts from

- accessions of *Amomum compactum* fruits. *Ann Agric Sci.* 2021;66(1):58-62. doi: 10.1016/j.aos.2021.04.001.
20. Purnomo D, Yulianti Putri A, Hisyam HM, Rimatunnisa R, Indriani DR, Adinda PR, et al. Effect of drying method on antioxidant activity and total flavonoid content of java tea crude drug (*Orthosiphon aristatus*). *Res Chem Eng.* 2023;2(1):29-33.
 21. Popoola OK. Phytotherapeutic potentials of *Synedrella nodiflora*: *In vitro* quantification of phytochemical constituents, anti-oxidant capacities and skin enzymes inhibiting activities. *Am J Biomed Sci Res.* 2020;10(6):543-50.
 22. Chen F, Huang G, Yang Z, Hou Y. Antioxidant activity of *Momordica charantia* polysaccharide and its derivatives. *Int J Biol Macromol.* 2019;138:673-80. doi: 10.1016/j.ijbiomac.2019.07.129, PMID 31344411.
 23. Ramasamy A, Anandakumar K, Kathiresan K. *In vitro* antioxidant potential and acetylcholinesterase inhibitory effect of *Ficus benghalensis* aerial root extract. *Afr Health Sci.* 2022;22(4):291-9. doi: 10.4314/ahs.v22i4.34, PMID 37092053.
 24. Le HT, Park J, Ha J, Kusumaningrum S, Paik JH, Cho S. *Synedrella nodiflora* (Linn.) Gaertn. inhibits inflammatory responses through the regulation of Syk in RAW 264.7 macrophages. *Exp Ther Med.* 2020;20(2):1153-62. doi: 10.3892/etm.2020.8750, PMID 32765663.
 25. HDT M. *In vitro* Anti-Inflammatory Egg Albumin Denaturation Assay: An Enhanced Approach. *J Nat Ayurvedic. Med.* 2023;7(3):1-6.
 26. Mans RA, Fripuron P, Djotaroeno M, Misser VS, Pawirodihardjo J. *In vitro* anti-inflammatory and antioxidant activities as well as phytochemical content of the fresh stem juice from *Montrichardia arborescens* Schott (Araceae). *Pharmacogn J.* 2022;14(4):296-304. doi: 10.5530/pj.2022.14.99.
 27. Anokwah D, Kwatia EA, Amponsah IK, Jibira Y, Harley BK, Ameyaw EO, et al. Evaluation of the anti-inflammatory and antioxidant potential of the stem bark extract and some constituents of *Aidia genipiflora* (DC.) dandy (Rubiaceae). *Heliyon.* 2022;8(8):e10082. doi: 10.1016/j.heliyon.2022.e10082, PMID 36042726.
 28. Fronza M, Heinzmann B, Hamburger M, Laufer S, Merfort I. Determination of the wound healing effect of *Calendula* extracts using the scratch assay with 3T3 fibroblasts. *J Ethnopharmacol.* 2009;126(3):463-7. doi: 10.1016/j.jep.2009.09.014, PMID 19781615.
 29. Bolla SR, Mohammed Al-Subaie A, Yousuf Al-Jindan R, Papayya Balakrishna J, Kanchi Ravi P, Veeraraghavan VP, et al. *In vitro* wound healing potency of methanolic leaf extract of *Aristolochia saccata* is possibly mediated by its stimulatory effect on collagen-1 expression. *Heliyon.* 2019;5(5):e01648. doi: 10.1016/j.heliyon.2019.e01648, PMID 31193473.
 30. Saleem U, Khalid S, Zaib S, Anwar F, Akhtar MF, Hussain L, et al. Wound Healing Potential and *in silico* Appraisal of *Convolvulus arvensis* L. Methanolic extract. *BioMed Res Int.* 2022;1373160. doi: 10.1155/2022/1373160, PMID 36467883.
 31. Governa P, Carullo G, Biagi M, Rago V, Aiello F. Evaluation of the *in vitro* wound-healing activity of Calabrian honeys. *Antioxidants (Basel).* 2019;8(2):36. doi: 10.3390/antiox8020036, PMID 30736314.
 32. Dhawal PP, Gharpure M, Joshi MS, Khan RR, Barve SS. Wound healing potential of Vakeri fortified Kampillakadi Taila. *J Ayurveda Integr Med.* 2023;14(3):100721. doi: 10.1016/j.jaim.2023.100721, PMID 37245340.
 33. Adil M, Filimban FZ, Ambrin QA, Quddoos AA, Sher AA, Naseer M. Phytochemical screening, HPLC analysis, antimicrobial and antioxidant effect of *Euphorbia parviflora* L. (*Euphorbiaceae* Juss.). *Sci Rep.* 2024;14(1):5627. doi: 10.1038/s41598-024-55905-w, PMID 38454096.
 34. Adedokun KA, Imodoye SO, Bello IO, Lanahun AA, Bello IO. Therapeutic potentials of medicinal plants and significance of computational tools in anti-cancer drug discovery. In: *Phytochemistry, computational tools and databases in drug discovery*. Amsterdam: Elsevier; 2023. p. 393-455. doi: 10.1016/B978-0-323-90593-0.00017-4.
 35. Dias MC, Pinto DC, Silva AM. Plant flavonoids: chemical characteristics and biological activity. *Molecules.* 2021;26(17):5377. doi: 10.3390/molecules26175377, PMID 34500810.
 36. Zulkefli N, Che Zahari CN, Sayuti NH, Kamarudin AA, Saad N, Hamezah HS, et al. Flavonoids as potential wound-healing molecules: emphasis on pathways perspective. *Int J Mol Sci.* 2023;24(5):4607. doi: 10.3390/ijms24054607, PMID 36902038.
 37. Mortada EM. Evidence-based complementary and alternative medicine in current medical practice. *Cureus.* 2024;16(1):e52041. doi: 10.7759/cureus.52041, PMID 38344508.
 38. Zhang Y, Cai P, Cheng G, Zhang Y. A brief review of phenolic compounds identified from plants: their extraction, analysis and biological activity. *Nat Prod Commun.* 2022;17(1):1934578X2110697. doi: 10.1177/1934578X211069721.
 39. Sanvee S, Simalou O, Tchani GW, Kagnou H, Bakoma B, Metowogo K, et al. Antidiabetic activity of tannin fraction of *Bridelia ferruginea* (Benth) leaf extract on fructose-induced diabetic mice. *J HerbMed Pharmacol.* 2020;10(1):68-74. doi: 10.34172/jhp.2021.06.
 40. Sravan Kumar S, Manoj P, Giridhar P. Fourier transform infrared spectroscopy (FTIR) analysis, chlorophyll content and antioxidant properties of native and defatted foliage of green leafy vegetables. *J Food Sci Technol.* 2015;52(12):8131-9. doi: 10.1007/s13197-015-1959-0, PMID 26604386.
 41. YENER İ. Trace element analysis in some plants species by inductively coupled plasma optical emission spectrometry (ICP-OES). *Iğdir Univ Fen Bilim Enstitüsü Derg.* 2019;9(3):1492-502. doi: 10.21597/jist.517739.
 42. NCBI. Trace elements-diet and health-NCBI bookshelf. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK218751/>.
 43. Bonaventura P, Benedetti G, Albarède F, Miossec P. Zinc and its role in immunity and inflammation. *Autoimmun Rev.* 2015;14(4):277-85. doi: 10.1016/j.autrev.2014.11.008, PMID 25462582.
 44. Salvo J, Sandoval C. Role of copper nanoparticles in wound healing for chronic wounds: literature review. *Burns Trauma.* 2022;10:tkab047. doi: 10.1093/burnst/tkab047, PMID 35071652.
 45. Kresovich JK, Bulka CM, Joyce BT, Vokonas PS, Schwartz J, Baccarelli AA, et al. The inflammatory potential of dietary manganese in a cohort of elderly men. *Biol Trace Elem Res.* 2018;183(1):49-57. doi: 10.1007/s12011-017-1127-7, PMID 28822065.
 46. Khaliq H, Juming Z, Ke-Mei P. The physiological role of boron on health. *Biol Trace Elem Res.* 2018;186(1):31-51. doi: 10.1007/s12011-018-1284-3, PMID 29546541.
 47. Hatami T, Emami SA, Miraghaee SS, Mojarab M. Total phenolic contents and antioxidant activities of different extracts and fractions from the aerial parts of *Artemisia biennis* Willd. *Iran J Pharm Res.* 2014;13(2):551-9. PMID 25237350.
 48. Saeed N, Khan MR, Shabbir M. Antioxidant activity, total phenolic and total flavonoid contents of whole plant extracts *Torilis leptophylla* L. *BMC Complement Altern Med.* 2012;12:221. doi: 10.1186/1472-6882-12-221, PMID 23153304.
 49. Aryal S, Baniya MK, Danekhu K, Kunwar P, Gurung R, Koirala N. Total phenolic content, flavonoid content and antioxidant potential of wild vegetables from Western Nepal. *Plants (Basel).* 2019;8(4):96. doi: 10.3390/plants8040096, PMID 30978964.
 50. Dharmadeva S, Galgamuwa LS, Prasadine C, Kumarasinghe N. *In vitro* anti-inflammatory activity of *Ficus racemosa* L. bark using albumin denaturation method. *Ayu.* 2018;39(4):239-42. doi: 10.4103/ayu.AYU_27_18, PMID 31367147.
 51. Tyavambiza C, Meyer M, Wusu AD, Madiehe AM, Meyer S. The antioxidant and *in vitro* wound healing activity of *Cotyledon orbiculata* aqueous extract and the synthesized biogenic silver nanoparticles. *Int J Mol Sci.* 2022;23(24):16094. doi: 10.3390/ijms232416094, PMID 36555732.

Cite this article: Koodathil J, Ramadass L, Jayachandran S, Sekar J. Phyto-Analytical Studies and *in vitro* Wound Healing Activity of *Synedrella nodiflora* (L.) Gaertn Leaf Extract. *Pharmacog Res.* 2025;17(2):x-x.