

Phytochemical Screening and Gas Chromatography-Mass Spectrometry Analysis of Ethanol Extract of *Scambiosia columbabria* L.

Idowu Jonas Sagbo, Ayuk Elizabeth Orock¹, Elizabeth Kola, Wilfred Otang-Mbeng

School of Biology and Environmental Sciences, University of Mpumalanga, Private Bag X11283, Mbombela, 1200, South Africa, ¹Department of Environmental Science, University of Buea, P. O. Box 63 Buea, South West Region, Cameroon

ABSTRACT


Background: *Scambiosia columbabria* L. is commonly known as the wild scabious and is used in South Africa to treat various ailments such as wound bruises, painful menstruation, heartburn, and colic. Despite its extensive traditional usage, the bioactive constituents of this plant are yet to be explored. **Objective:** The present study was carried out to identify the bioactive constituents of the ethanol extract of *S. columbabria* by phytochemical screening and gas chromatography-mass spectrometry (GC-MS). **Materials and Methods:** Sixty grams of the powdered leaves was sequentially extracted by ethanol and later tested for preliminary phytochemical screening and further subjected to GC-MS analysis. **Results:** The results showed the presence of alkaloids, flavonoids, phenolics, tannins, steroids, terpenoids, and saponins in the extract. The GC-MS analysis revealed the presence of 16 major compounds, with flavonoids (40.84%) being the most represented chemical class. **Conclusion:** The findings indicated that the plant possesses compounds with biological activities and therefore justifies its traditional usage in the treatment of skin and other diseases.

Key words: Biological activity, diseases, gas chromatography-mass spectrometry, phytochemical screening, *Scambiosia columbabria*

SUMMARY

Scambiosia columbabria L. is commonly known as the wild scabious and is used in South Africa to treat various ailments such as wound bruises, painful menstruation, heartburn, and colic. The result generated from the study reveals the presence of various bioactive compounds which are

known to exhibit various biological activities and therefore validates the reports of therapeutic importance of this plant in the treatment of skin and other diseases, and it may therefore be recommended as plant of pharmaceutical importance.



Abbreviations Used: GC-MS: Gas chromatography-mass spectrometry, MSD: Mass selective detector, WHO: World Health Organization.

Correspondence:

Dr. Idowu Jonas Sagbo,
School of Biology and Environmental Sciences,
University of Mpumalanga, Private Bag X11283,
Mbombela, 1200, South Africa.
E-mail: jonas.sagbo@ump.ac.za
DOI: 10.4103/pr.pr_59_19

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Recently, the use of plant products in ethnomedicine is increasing globally. The World Health Organization has also pointed out that about 80% of the world's population rely heavily on herbal medicinal products for the treatment of various health challenges.^[1,2] These products are being prepared as decoction, infusions, extracts, or pastes with easily accessible liquids such as alcohol, water, and milk. In South Africa, the use of herbs to treat various illnesses continues to expand rapidly across all provinces, and studies have also indicated that South Africa hosts a variety of about 30,000 plant species which are used for treating numerous ailments such as skin, cancer, diabetes, tuberculosis, and hypertension.^[3] However, the bioactive constituents responsible for the therapeutic activity of many South African medicinal plants have not been scientifically elucidated, and such is the case of *Scambiosia columbabria* which is investigated in the current study.

S. columbabria L. is a perennial herb that grows to about 1 m in height, with annual branches developing from persistent fleshy roots.^[4] It belongs to the family Caprifoliaceae (formerly Dipasacaceae). The plant is locally called wild scabious (English); bitterwortel (Afrikaans); makgha (Xhosa); ibheka (Zulu); and moholungoane (South Sotho). *S. columbaria* occurs mostly in grasslands and rocky slopes, is widely

distributed throughout South Africa, and is particularly common in the Western Cape. Conventionally, the dried roasted roots of the plant are made into a wound-healing ointment to treat wound bruises, and the roots are used to prepare a pleasant smelling baby powder.^[5] In addition, the root and leaves are also used to treat painful menstruation, heartburn, and colic. Despite the extensive traditional usage of this plant, to date, no published *in-vitro* or *in-vivo* studies are available in literature. Some of the compounds previously identified from the rootstock of *S. columbabria* include loganin and sweroside.^[4] However, there is still a lack of comprehensive studies with regard to the active constituents of this plant. Hence, the present study was carried out to identify the bioactive constituents of the ethanol extract of *S. columbabria* by phytochemical screening and gas

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Sagbo IJ, Orock AE, Kola E, Otang-Mbeng W. Phytochemical screening and gas chromatography-mass spectrometry analysis of ethanol extract of *Scambiosia columbabria* L. Phcog Res 2020;12:35-9.




Figure 1: Chemical structure of identified compounds from *Scambiosa columbabria*. Tetraethyl silicate (1) Eucalyptol (2), Benzene, 1,4-dichloro- (3), 2-Naphthalenemethanol, 1,2,3,4,4a, 5,6,7-octahydro-.alpha., alpha., 4a, 8-tetramethyl-, (2R-cis)- (4), 2-Naphthalenemethanol, decahydro-.alpha., alpha,4a-trimethyl-8-methylene-, [2R-(2.alpha.,4.alpha.,8.alpha.)]- (5), Phytol (6), Aflatoxin B2 (7), Pregna-5-ene-3,20-dione (8), Podocarpa-8,11,13-trien-3-one, 14-isopropyl-13-methoxy-(9)H-1-Benzopyran-4-one,5-hydroxy-2-(4-hydroxyphenyl)-3,7-dimethoxy-(10), Batilol (11), 1-Methyl-3-[(1R)-1-phenylpropyl]thiourea(12),Aprobarbital(13),Acetamide,N-benzyl-2-(3-cyano-4,6-dimethylpyridin-2-ylsulfanyl)-N-ethyl-(14), 3,5-Dihydroxy-6,7,8-trimethoxyflavone (15), 4-(para-Chlorobenzyl)-pyridine (16)




Figure 2: Gas chromatography-mass spectrometry chromatogram of ethanolic extract of *Scambiosa columbabria* leaves

Table 1: Qualitative screening of phytochemical components present in the ethanol extract of *Scambiosa columbabria*

Phytochemical test	Name of the test	Ethanol extract of <i>Scambiosa columbabria</i>
Alkaloids	Mayer's test	+
Flavonoids	Alkaline reagent test	+
Phenols	Ferric chloride test, lead acetate test	+
Tannins	Gelatin test	+
Steroids	Liebermann-burchard's test	+
Terpenoids	Copper acetate test	+
Saponin	Frothing test	+

+: Present

chromatography-mass spectrometry (GC-MS) with the view to justify its ethnobotanical usage.

MATERIALS AND METHODS

Plant collection

The whole plant material together with the leaves of *S. columbabria* was collected from Alice, Eastern Cape province of South Africa, during the month of January, 2019, at 10:00 AM. The collected plant was authenticated by a botanist at the Giffen Herbarium of the University of Fort Hare, where voucher specimens are kept. The leaves of *S. columbabria* were detached from the rest of the plant after which they were washed with clean tap water thrice to eradicate debris and later oven-dried to a constant weight at 30°C for 1 week. Thereafter, the dried leaves were then milled to a homogenous powder using an electric blender (Polymix PX-MFC 90D, KINEMATICA AG, Luzern, Switzerland).

Plant extraction

About 60 g of the powdered leaves was extracted in 1000 ml of ethanol (99.99%) and placed on an orbital shaker (Labcon laboratory service [Pty], South Africa) for 24 h. The extract was later filtered using a Buchner funnel and Whatman No. 1 filter paper. The resulting filtrate was concentrated to dryness using a rotary evaporator (Heidolph Laborata 4000, Heidolph instruments, GmbH and Co, Germany). The final extract was weighed to determine the yield (4.54%) and then stored at 4°C in a refrigerator for further studies.

Table 2: The major phytochemicals detected in the ethanolic leaf extract of *Scambiosia columbabra* by gas chromatography-mass spectrometry analysis

Retention time (min)	Compound name	Molecular formula	Molecular weight	Area (%)
11.602	Tetraethyl silicate	C ₈ H ₂₀ O ₄ Si	208	0.79
13.055	Eucalyptol	C ₁₀ H ₁₈ O	154	1.11
13.17	Benzene, 1,4-dichloro-	C ₆ H ₄ Cl ₂	147	2.24
38.548	2-Naphthalenemethanol, 1,2,3,4,4a, 5,6,7-octahydro-.alpha.,.alpha.,4a, 8-tetramethyl-, (2R-cis)-	C ₁₅ H ₂₆ O	222	4.45
39.209	2-Naphthalenemethanol, decahydro-.alpha.,.alpha.,4a-trimethyl-8-methylene-, [2R-(2 α ,4 α ,8 α)]-	C ₁₅ H ₂₆ O	222	3.66
54.639	Phytol	C ₂₀ H ₄₀ O	296	0.54
56.005	Aflatoxin B2	C ₁₇ H ₁₄ O ₆	314	10.45
57.161	Pregn-5-ene-3,20-dione	C ₂₁ H ₃₀ O ₂	314	0.35
58.546	Podocarpa-8,11,13-trien-3-one, 14-isopropyl-13-methoxy-	C ₂₁ H ₃₀ O	314	13.71
59.325	4H-1-Benzopyran-4-one, 5-hydroxy-2-(4-hydroxyphenyl)-3,7-dimethoxy-	C ₁₇ H ₁₄ O ₆	314	0.68
62.286	Batilol	C ₂₁ H ₄₄ O ₃	344	6.86
62.592	1-Methyl-3-[(1R)-1-phenylpropyl] thiourea	C ₁₁ H ₁₆ N ₂ S	208	1.71
63.602	Aprobarbital	C ₁₀ H ₁₄ N ₂ O ₃	210	2.1
65.532	Acetamide, N-benzyl-2-(3-cyano-4,6-dimethylpyridin-2-ylsulfanyl)-N-ethyl-	C ₁₉ H ₂₁ N ₃ OS	339	1.73
67.176	3,5-Dihydroxy-6,7,8-trimethoxyflavone	C ₁₈ H ₁₆ O ₇	344	40.84
67.403	4-(para-Chlorobenzyl)-pyridine	C ₁₂ H ₁₀ ClN	203	8.78
Total identified components (%)				100

Table 3: Nature of the compounds identified in the extract of *Scambiosia columbabra*

Compound name	Nature of compound
Tetraethyl silicate	Ester
Eucalyptol	Monoterpenoid
Benzene, 1,4-dichloro-	Phenol
2-Naphthalenemethanol, 1,2,3,4,4a, 5,6,7-octahydro-.alpha.,.alpha.,4a, 8-tetramethyl-, (2R-cis)-	NA
2-Naphthalenemethanol, decahydro-.alpha.,.alpha.,4a-trimethyl-8-methylene-, [2R-(2 α ,4 α ,8 α)]-	NA
Phytol	Diterpenes
Aflatoxin B2	Coumarins
Pregn-5-ene-3,20-dione	Steroid
Podocarpa-8,11,13-trien-3-one, 14-isopropyl-13-methoxy-	Steroid
4H-1-Benzopyran-4-one, 5-hydroxy-2-(4-hydroxyphenyl)-3,7-dimethoxy-	NA
Batilol	Alkyl glycerol
1-Methyl-3-[(1R)-1-phenylpropyl] thiourea	NA
Aprobarbital	Diazine
Acetamide, N-benzyl-2-(3-cyano-4,6-dimethylpyridin-2-ylsulfanyl)-N-ethyl-	NA
3,5-Dihydroxy-6,7,8-trimethoxyflavone	Flavonoid
4-(para-Chlorobenzyl)-pyridine	Pyridine

N/A: Not available in database

Table 4: Class composition of the compounds identified in the extract of *Scambiosia columbabra*

Class of compound	<i>Scambiosia columbabra</i>
Alkyl glycerol	6.86
Coumarins	10.45
Diterpenes	0.54
Diazines	2.1
Esters	0.79
Flavonoids	40.84
Monoterpenes	1.11
Phenols	2.24
Steroids	14.06
Others	21.01
Total grouped components (%)	100

Table 5: Reported biological activity of some compounds identified in the extract of *Scambiosia columbabra*

Name of compound	Reported biological activity	Reference
Eucalyptol	Anti-inflammatory, antioxidant and antibacterial	[37,39,40]
Benzene, 1,4-dichloro-	Deodorant	[32]
Phytol	Cytotoxic, antinociceptive, antioxidant, antimicrobial, anti-inflammatory, anticancer, diuretic	[37,41]
Pregn-5-ene-3,20-dione	Antibacterial and antilarva	[36]
Batilol	It is used in cosmetics as a stabilising ingredient and skin-conditioning agent	[37]
Aprobarbital	anticonvulsant properties and used for the treatment of insomnia	[32]
3,5-Dihydroxy-6,7,8-trimethoxyflavone	Antifungal	[38]

Phytochemical screening

The presence or absence of phytochemical constituents such as alkaloids, flavonoids, phenolics, tannins, steroids, terpenoids, and saponins was screened using standard qualitative procedures.^[6-8]

Gas chromatography-mass spectrometry analysis

The GC-MS analysis of the ethanol extract of *S. columbabria* was carried out using an Agilent 7890 GC system equipped with an Agilent 5977A Mass selective detector system (Chematrix [Pty] Ltd., Agilent Technologies, DE, Germany) with a Zebron-5MS (cross-linked 5% – phenyl methyl polysiloxane) column (HP-5 fused silica 30 m × 0.320 mm × 0.250 µm film thickness). The instrument was operated at injection temperature (250°C), ion source temperature (280°C), and pressure (48.745 kpa). The carrier gas used was GC-grade helium (99.999% purity) at a flow rate of 36.262 cm/s, and about 1 µL sample injections was used. The oven temperature was initially started from 40°C (held for 1 min) and then ramped to 240°C at 3°C/min.

Identification of components

Identification of components was conducted using the National Institute of Standard and Technology 11 library source. The mass spectrum of individual unknown compound was achieved by comparing their retention times with those of the known compound in the software database library.

RESULTS AND DISCUSSION

To investigate the importance of any medicinal plant, the initial or first step is to screen for its phytochemicals, as it gives a broad knowledge with respect to the nature of the compounds present in it. In the present study, the ethanolic leaf extracts of *S. columbabria* were preliminarily screened for the phytochemicals.

The extract shows the presence of alkaloids, flavonoids, phenolics, tannins, steroids, terpenoids, and saponins [Table 1]. These observed phytochemicals have been reported to possess a broad range of activities.^[9-11] Alkaloids present in the extract of *S. columbabria* have been reported to possess anticonvulsant,^[12] hypotensive,^[13] antimicrobial,^[12] antiprotozoal,^[14] and antimalarial activities.^[15] In addition, studies have also shown that plants with alkaloids may have hypoglycemic effect via insulin-releasing mechanism and insulin-mimicking activity, thus improving postprandial hyperglycemia.^[9] The presence of flavonoids in plant extract has also been reported to play an important role in plant biochemistry and physiology as enzyme inhibitor, antioxidants,^[16] anti-diabetic,^[17-19] anti-inflammatory,^[20] antimicrobial,^[21-23] antifungal,^[21] and anticancer activities, thereby protecting the cells against all stages of carcinogenesis.^[10] Furthermore, available report on phenolic compounds has also been demonstrated for their usefulness to prevent oxidative damage in cells, thereby acting as a free radical terminator.^[24] Tanins, steroids, and terpenoids present in the extract have been responsible for anti-inflammatory, antioxidant, antibacterial, and hypoglycemic activities.^[25,26] It can be observed from this study that the presence of alkaloids, flavonoids, phenolics, tannins, steroids, terpenoids, and saponins in the extract of *S. columbabria* leaves is in support with those reported from previous studies, indicating the presence of these phytochemicals in the plant extracts.^[27-29] Therefore, these phytochemicals observed in the ethanol leaf extract of *S. columbabria* are an indication that this plant plays a vital role as a dietary antioxidant, and it can be used to develop new compounds for health benefit.

The results of the GC-MS analysis of *S. columbabria* revealed the presence of sixteen compounds [Figures 1 and 2], representing 100%

of the total identified components [Table 2]. The detected compounds class were mainly composed of alkyl glycerol, coumarins, diterpenes, esters, flavonoids, monoterpenes, steroids, phenols, and others [Table 3]. The most represented chemical class were flavonoids (40.84%), steroids (14.06%), coumarins (10.45%), alkyl glycerols (6.86%), phenols (2.24%), monoterpenes (1.11%), ester (0.79%), and diterpenes (0.54%) [Table 4]. However, studies have indicated that compounds mirror the biological activities of plant from which it is extracted. Among the compounds identified, eucalyptol which is a monoterpene found in the extract of *S. columbabria* was reported to control asthma via anti-inflammatory cytokine inhibition [Table 5].^[30] In addition to anti-inflammatory activity, eucalyptol was also reported to be frequently used in the manufacturing of cosmetics, thereby increasing the percutaneous penetration of drugs as aromatherapy agent.^[31] Benzene, 1, 4-dichloro-, one of the main components found in the extract of *S. columbabria*, was reported being used as a deodorant.^[32] Phytol, a key acyclic diterpene class of compounds, identified in the extract, was also reported to exhibit antibacterial activity against *Staphylococcus aureus* and *Escherichia coli* by causing enormous damage to the bacterial cell membrane.^[33] Pejín *et al.*^[34] also reported strong antimicrobial activity of phytol by microdilution method against eight bacterial and eight fungal strains. In addition to its antimicrobial activity, phytol was also reported to exhibit no remarkable toxicity when treated with mouse skin cells.^[35] Pregn-5-ene-3, 20-dione, a steroid detected in the extract, has been indicated to possess potent antibacterial activity against 15 marine bacterial strains.^[36] Batilol, found in the ethanol extract of *S. columbabria*, was reportedly used in cosmetics as a stabilizing ingredient and skin-conditioning agent,^[37] while 3,5-Dihydroxy-6,7,8-trimethoxyflavone, a flavonoid, detected in higher amount (40.84%) in the extract, was previously reported to show moderate antifungal activity against *Candida albicans* with an minimum inhibitory concentration value of 312 µg/ML.^[38] However, to the best of our knowledge, this is the first study to identify the bioactive constituents of *S. columbabria* by phytochemical screening and GC-MS. Therefore, it could be concluded from this study that ethanol extract of *S. columbabria* composed of various bioactive compounds which are known to exhibit various biological activities and may therefore be recommended as a plant of pharmaceutical importance.

CONCLUSION

The study shows that ethanol leaf extracts of *S. columbabria* contained some phytoconstituents which are pharmacologically important and therefore could represent potential source of lead molecules with pharmacological activities for the development of new novel drug or a leading compound for the treatment of skin and other diseases. In addition, the presence of these compounds with biological activities justifies the traditional usage of *S. columbabria* for the treatment of skin and other diseases. Hence, further studies are needed to isolate individual bioactive compounds responsible for its therapeutic activity and the elucidation of their mechanism(s) of action.

Acknowledgements

The authors would like to express their gratitude to the National Research Foundation (Grant no: 105161) and University of Mpumalanga, for financial support to conduct this research.

Financial support and sponsorship

This study was financially supported by National Research Foundation (grant no: 105161).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- World Health Organization. WHO Guidelines on Safety Monitoring of Herbal Medicines in Pharmacovigilance Systems. Geneva, Switzerland: World Health Organization; 2004.
- Ekor M. The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol* 2014;4:177.
- Louw CA, Regnier TJ, Korsten L. Medicinal bulbous plants of South Africa and their traditional relevance in the control of infectious diseases. *J Ethnopharmacol* 2002;82:147-54.
- van Wyk BE, van Oudtshoorn B, Gericke N. Medicinal Plants of South Africa. Pretoria: Briza Publications; 2009.
- Von Koenen E. Medicinal, Poisonous and Edible Plants in Namibia. Windhoek: Klaus Hess Publishers; 2001.
- Ajayi GO, Olagunju JA, Ademuyiwa O, Martins OC. Gas chromatography-mass spectrometry analysis and phytochemical screening of ethanolic root extract of *Plumbago zeylanica*, Linn. *J Med Plants Res* 2011;9:1756-61.
- Nwilo BI, Uwakwe AA, Akaninwo JO. Phytochemical screening and GC-FID analysis of ethanolic extract of root bark of *Salacia nitida* L. Benth. *J Med Plants Stud* 2016;4:283-7.
- Swamy MK, Sinniah UR, Akhtar MS. *In vitro* pharmacological activities and GC-MS analysis of different solvent extracts of *Lantana camara* leaves collected from tropical region of Malaysia. *Evid Based Complement Alternat Med* 2015;2015:506413.
- Patel MB, Mishra S. Hypoglycemic activity of alkaloidal fraction of *Tinospora cordifolia*. *Phytomedicine* 2011;18:1045-52.
- Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and extraction: A review. *Int Pharm* 2011;1:98-106.
- Devendran G, Balasubramanian U. Qualitative phytochemical screening and GC – MS analysis of *Ocimum sanctum* L. leaves. *Asian J Plant Sci Res* 2011;1:44-8.
- Singh H, Kappor V. Investigation of *Strychnos* Spp. V1. Pharmacological studies of alkaloids of *Strychnos potatorum* seeds. *Planta Med* 1980;38:133-7.
- Ali MM, Ghatak BJ. Pharmacological investigation of an alkaloid fraction isolated from *Strychnos potatorum* Linn. *Indian J Exp Biol* 1975;13:163-7.
- Frédérich M, Jacquier MJ, Thépenier P, De Mol P, Tits M, Philippe G, *et al.* Antiplasmodial activity of alkaloids from various *Strychnos* species. *J Nat Prod* 2002;65:1381-6.
- Wadood A, Ghufuran M, Jamal SB, Naeem M, Khan A, Ghaffar R, *et al.* Phytochemical analysis of medicinal plants occurring in local area of Mardan. *Biochem Anal Biochem* 2013;2:144.
- Yamamoto Y, Gaynor RB. Therapeutic potential of inhibition of the NF-kappaB pathway in the treatment of inflammation and cancer. *J Clin Invest* 2001;107:135-42.
- Tanko Y, Yerima M, Mahdi MA, Yaro AH, Musa KY, Mohammed A. Hypoglycemic activity of methanolic stem bark of *Adansoniadigitata* extract on blood glucose levels of streptozocin-induced diabetic wistar rats. *Int J Appl Res Nat Prod* 2008;1:32-6.
- Moses AG, Maobe Gitu L, Erastus G, Rotich H. Phytochemical analysis of phenol and flavonoid in eight selected medicinal herbs used for the treatment of diabetes, malaria and pneumonia in Kisii, Kenya. *Acad J Cancer Res* 2012;5:31-9.
- Sharma R. Preliminary phytochemical screening of some indigenous medicinal plant leaves extract in regulation of antidiabetic activity. *Sci Res Rep* 2012;2:307-10.
- Prabhu VV, Guruvayoorappan C. Phytochemical screening of methanolic extract of mangrove *Avicennia marina* (Forssk.) Vierh *Der Pharm Sin* 2012;3:64-70.
- Lima AL, Parial R, Das M, Das AK. Phytochemical and pharmacological studies of ethanolic extract from the leaf of mangrove plant *Phoenix paludosa* Roxb. *Malays J Pharm Sci* 2010;8:59-69.
- Gawali P, Jadhav BL. Antioxidant activity and antioxidant phytochemical analysis of mangrove species *Sonneratia alba* and *Bruguiera cylindrica*. *Asian J Microbiol Biotechnol Environ Sci* 2011;13:257-61.
- Nurdiani R, Firdaus M, Prihanto AA. Phytochemical screening and antibacterial activity of methanol extract of mangrove plant *Rhizophora mucronata* from Porong River Estuary *J Basic Sci Technol* 2012;1:27-9.
- Block G. The data support a role for antioxidants in reducing cancer risk. *Nutr Rev* 1992;50:207-13.
- Akindele AJ, Adeyemi OO. Antiinflammatory activity of the aqueous leaf extract of *Byrsocarpus coccineus*. *Fitoterapia* 2007;78:25-8.
- likay O, Esra K, Bilge S, Erdem Y. Appraisal of anti-inflammatory potential of the clubmoss, *Lycopodium clavatum* L. *J Ethnopharmacol* 2007;109:146-50.
- Starlin T, Raj CA, Ragavendran P, Gopalakrishnan VK. Phytochemical screening, functional group and elemental analysis of *Tylophora pauciflora* Wight and Arn. *Int Res J Pharm* 2012;3:180-3.
- Revathi PT, Jeyaseelansenthinath T, Thirumalaikolundhusubramaian P. Preliminary phytochemical screening and GC-MS analysis of mangrove plant *Bruguiera Clinidrica* (Rhizho) L. *Int J Pharmacogn Phytochem Res* 2014;6:15.
- Tyagi T, Agarwal M. Phytochemical screening and GC-MS analysis of bioactive constituents in the ethanolic extract of *Pistia stratiotes* L. and *Eichhornia crassipes* (Mart) solms. *J Pharmacogn Phytochem* 2017;6:195-206.
- Juergens UR, Stöber M, Schmidt-Schilling L, Kleuver T, Vetter H. Antiinflammatory effects of euclyptol (1,8-cineole) in bronchial asthma: Inhibition of arachidonic acid metabolism in human blood monocytes *ex vivo*. *Eur J Med Res* 1998;3:407-12.
- Dogan H, Tasman F, Cehreli ZC. Effect of gutta-percha solvents at different temperatures on the calcium, phosphorus and magnesium levels of human root dentin. *J Oral Rehabil* 2001;28:792-6.
- Wishart DS, Feunang YD, Marcu A, Guo AC, Liang K. HMDB 4.0 — The Human Metabolome Database for 2018 *Nucleic Acids Res* 2018;46:D608–17. 29140435, 2018.
- Nandagopalan V, Johnson GM, Doss A. GC-MS analysis of bioactive components of the methanol extract of *Hibiscus tiliaceus* Linn. *Asian J Plant Sci Res* 2015;5:6-10.
- Pejin B, Savic A, Sokovic M, Glamoclija J, Ciric A, Nikolic M, *et al.* Further *in vitro* evaluation of antiradical and antimicrobial activities of phytol. *Nat Prod Res* 2014;28:372-6.
- Ghaneian MT, Hassan HE, Jebali A, Hekmatimoghaddam S, Mahmoudi M. Antimicrobial activity, toxicity and stability of phytol as a novel surface disinfectant. *Environ Heal Eng Manag J* 2015;2:1.
- Faheem A, Yam WWS, Koay YC. Chemical constituents and biological activities of the genus *Subergorgia*. *Pharmacogn Rev* 2012;6:78-80.
- Duke J. Phytochemical and ethnobotanical database. Available from: <https://phytochem.nal.usda.gov/phytochem/search>. [Last accessed on 2019 Mar 12].
- Doğan H, Taşman F, Cehreli ZC. Effect of gutta-percha solvents at different temperatures on the calcium, phosphorus and magnesium levels of human root dentin. *J Oral Rehabil* 2001;28:792-6.
- Pattnaik S, Subramanyam VR, Bapaji M, Kole CR. Antibacterial and antifungal activity of aromatic constituents of essential oils. *Microbios* 1997;89:39-46.
- Karlovic Z, Anic I, Miletic I, Prpic-Mehicic G, Pezelj-Ribaric S, Marsan T. Antibacterial Activity of Halothane Eucalyptol and Orange Oil. *Acta Stomat Croat* 2000;22:307-9.
- Prakasia PP, Nair A. Chemical fingerprint of essential oil components from fresh leaves of *Glycosmis pentaphylla* (Retz.) Correa. *Pharma Innov J* 2015;3:50–6.