

GC-MS Analysis of Bioactive Compounds Present in the Different Extracts of the Whole Plant *Impatiens minor* (DC.) Bennet

Akila Sukumaran^{1,*}, Ansa Mathew², Sebastin Varghese³, Ajithbabu Tharackal Kannankutty³

¹Department of Pharmacognosy, Malik Deenar College of Pharmacy, Seethangoli, Kasaragod, Kerala, INDIA.

²Department of Pharmacognosy, Dr. Joseph Mar Thoma Institute of Pharmaceutical Sciences and Research, Kayamkulam, Alappuzha, Kerala, INDIA.

³Department of Pharmaceutical Chemistry, Malik Deenar College of Pharmacy, Seethangoli, Kasaragod, Kerala, INDIA.

ABSTRACT

Background and Objectives: The systematic assessment of herbs utilized in the traditional medicine could provide more promising data regarding their therapeutic properties helping to address the increasing need for innovative treatments for infections and illnesses. Considering this perspective, *Impatiens minor*, an important medicinal plant distributed in the wet wastelands throughout India and also in the wet forests of Western Ghats was selected for our research. **Materials and Methods:** In this study, the plant *I. minor* in whole was collected, authenticated and subjected to shade drying for powdering which was done by mechanical grinder. Soxhlation technique was employed for the extraction of powdered plant material using different solvents to prepare the dried extract which was subjected to preliminary phytochemical evaluation and GC-MS analysis. **Results:** Presence of glycosides, phenolic compounds, flavonoids, terpenoids, sterols and carbohydrates was found in the preliminary phytochemical evaluation of different extracts. In the GC-MS analysis, presence of eight major compounds were found in the petroleum ether extract, while the chloroform extract showed the presence of six major compounds and twelve major compounds were found in the ethanol extract. **Conclusion:** It is worth noting that certain compounds identified in the GC-MS analysis have been documented to possess anti-oxidant, anti-inflammatory, and anticancer properties in previous studies. This study has the potential to contribute valuable insights for future in-depth investigations.

Keywords: *Impatiens minor* (DC.) Bennet, Extraction, Preliminary phytochemical evaluation, GC-MS analysis.

Correspondence:

Associate Prof. Akila Sukumaran

Department of Pharmacognosy,
Malik Deenar College of Pharmacy,
Seethangoli, Kasaragod, Kerala, INDIA.
Email: akhilasuku@yahoo.com

Received: 24-09-2024;

Revised: 04-11-2024;

Accepted: 06-12-2024.

INTRODUCTION

For centuries, nature has served as a rich reservoir of medicinal compounds, providing humanity with a vast array of healing agents. Throughout history, numerous modern pharmaceuticals have been developed by harnessing the power of natural sources. In fact, a significant proportion of these breakthroughs can be traced back to the traditional uses of these agents in ancient medicine.^[1] Nowadays, there is a resurgence in the fascination with conventional medicine and a growing need for additional medications derived from plants. The resurgence in the popularity of plant-based medications primarily stems from the prevailing notion that "green medicine" is secure and more reliable compared to expensive synthetic drugs, a significant number of which have

undesirable side effects.^[2] Phytochemicals play a crucial role in the management of several diseases including asthma, arthritis, cancer, etc. Numerous potent phytochemicals have been extracted and characterized from these green factories, leading to the development of several drugs with exceptional efficacy.^[3] In the recent decades, gas chromatography-mass spectrometry (GC-MS) has established firmly as a key technological tool for the characterization of phytochemicals.^[4] With this view, the plant *Impatiens minor* (DC.) Bennet (Family of *Balsaminaceae*) (Figure 1) was selected for the present study.

The genus *Impatiens* has around 1000 species of plants and new species are constantly discovered. Most of *Impatiens* species have a palaeotropical origin, however, there are few species endemic from Eurasia, North America, and Central America. Several plants of *Impatiens* genus have relevance for agriculture, ethno medicine and pharmacology.^[5-9] The *I. minor* is one among them commonly known in the name of lesser balsam or wild balsam. It is also known in the synonym of *Balsamina minor* DC and *I.*



DOI: 10.5530/pres.20251952

Copyright Information :

Copyright Author (s) 2025 Distributed under
Creative Commons CC-BY 4.0

Publishing Partner : Manuscript Technomedia. [www.mstechnomedia.com]

kleinii Wight and Arn. It is a small succulent annual herb endemic to peninsular India and distributed widely in Western Ghats, usually growing in the rainy season. Leaves are simple, opposite, ovate-lanceolate, serrate, and pubescent. Flowers are rose coloured, axillary, solitary, or in pairs. Fruits, pale green dehiscent capsules, contain small globose black seeds. It well grows in moist places, rock crevices, on roofs or walls, in shade, and also on tree trunks.^[10,11] The primary object of this study was to conduct an preliminary phytochemical screening of the different extracts of the whole plant *I. minor* and the characterization of active constituents by GC-MS analysis. The findings of this study would serve as a solid foundation for future investigations.

MATERIALS AND METHODS

Plant material-Collection and Identification

The whole plant of *Impatiens minor* (DC.) Bennet was collected from Seethangoli, a small village located nearer to Kasaragod town of Kerala, India. Mr. Biju P., Assistant Professor, Department of Botany, Government College, Vidyanagar, Kasaragod, Kerala, India identified and authenticated the collected material.

Plant material-Powdering and Extraction

The collected whole plant material was powdered and extracted in reference with previous literature.^[12-16] After 30 days of shade drying, the plant material was powdered with mechanical grinder and stored in airtight container for further study. The successive solvent extraction of the coarse powdered material (35 g) was done with the solvents (500 mL) of ascending order of polarity viz., petroleum ether, chloroform, and ethanol (70%) by soxhlet extraction procedure. Each extract was filtered, distilled off the solvent to obtain pure dried extract which were preserved for further studies.

Preliminary phytochemical screening

Preliminary phytochemical screening of the prepared extracts was carried out in accordance with the procedure in the published literatures.^[12-17]

Analysis of extracts by GC-MS

GC-MS analysis of petroleum ether, chloroform, and ethanol extract was carried out using a Perkin Elmer GC/MS Claurus 680/600 system provided with an Elite-5MS column (30.0 m, 0.25 mm ID, 250 μ m d_p). For detection in GC/MS, an electron ionization system with ionization energy of 70eV was used. Helium gas (99.999%) was employed as the carrier gas at a constant flow rate of 1 mL/min and an injection volume of 1 μ L was used (split ratio of 10:1). Injection port temperature was ensured as 250°C and ion-source temperature 230°C. The instrument was set at an initial temperature of 60°C and maintained this temperature for 2 min. At the end of this period, the oven temperature reached 300°C, at the rate of rise of 10°C/min, and maintained for 6 min. Mass spectra were taken at 70 eV; a scan interval of 0.2 sec and fragments from 50 to 600Da. Total GC running time was 32 min. The relative percentage amount of each component was calculated by comparing its average peak












Figure 1: *Impatiens minor* (DC.) Bennet.

Table 1: Preliminary qualitative phytochemical screening of the *Impatiens minor* extracts.

Sl. No.	Phytochemical	Petroleum ether extract	Chloroform extract	Ethanol extract
1.	Alkaloids	-	-	-
2.	Glycosides	-	-	+
3.	Phenolic compounds and Tannins	-	-	+
4.	Flavones and Flavonoids	-	-	+
5.	Carbohydrates	-	-	+
6.	Proteins and Amino acids	-	-	-
7.	Terpenoids	+	+	+
8.	Sterols	+	+	-
9.	Saponins	-	-	-
10.	Gum and mucilage	-	-	-

Table 2: Compound detected in the GC-MS scanning of petroleum ether extract of *I. minor*.

Sl. No.	Retention time	Name of the compound	Mol. Formula	Structure	Mol. Wt.	Peak Area %
1.	16.90	Z,Z-6,2 8-Heptatriactontadien-2-one	C ₃₇ H ₇₀ O		530	0.705
2.	16.98	4-methoxy-6-methyl- 6,7-dihydro-4H-Furo(3,2-C) pyran	C ₉ H ₁₂ O ₃		168	0.567
3.	19.60	Phytol	C ₂₀ H ₄₀ O		296	2.58
4.	23.17	1,2-Benzenedicarboxylic Acid, Mono (2-Ethylhexyl) Ester	C ₁₆ H ₂₂ O ₄		278	5.38
5.	26.52	Hentriacontane	C ₃₁ H ₆₄		436	4.78
6.	29.58	Cyclotrisiloxane, Hexamethyl-	C ₆ H ₁₈ O ₃ Si ₃		222	6.38
7.	30.19	2-oleanen-3-yl acetate,[3 alpha]	C ₃₂ H ₅₂ O ₂		468	19.92
8.	30.79	2R-acetoxymethyl- 1,3,3-trimethyl-4T- (3-methyl-2-buten-1-yl)-1t -cyclohexanol	C ₁₇ H ₃₀ O ₃		282	7.99

**Figure 2:** GC-MS investigation of petroleum ether extract of *I. minor*.










Figure 3: GC-MS scrutiny of chloroform extract of the *I. minor*.

Table 3: Compounds identified in the GC-MS assessment of chloroform extract of *I. minor*.

Sl.No.	Retention time	Name of the compound	Mol. Formula	Structure	Mol. Wt.	Peak Area %
1.	11.89	Benzene, 1,2-Dimethoxy-4-(2-Propenyl)	C ₁₁ H ₁₄ O ₂		178	3.73
2.	16.97	Sydnone, 3-(3,3-Dimethylbutyl)-	C ₈ H ₁₄ O ₂ N ₂		170	1.19
3.	19.58	Z,Z-6,2 8-Heptatriactontadien-2-one	C ₃₇ H ₇₀ O		530	11.64
4.	25.11	Hentriacontane	C ₃₁ H ₆₄		436	4.44
5.	26.50	Silane, Trichlorooctadecyl	C ₁₈ H ₃₇ C ₃ Si		386	5.16
6.	29.93	Cyclotrisiloxane, Hexamethyl-	C ₆ H ₁₈ O ₃ Si ₃		222	12.54

area to the total area. Turbo Mass Ver5.4.2 software was used to handle the mass spectra and chromatograms.^[18]

RESULTS

Preliminary Phytochemical Screening













Results of preliminary qualitative phytochemical screening of petroleum ether, chloroform and ethanol extract of whole plant of *I. minor* is shown in Table 1. Alkaloids was absent in the tested extracts. The ethanol extract showed the presence of glycosides, phenolic compounds, flavonoids and carbohydrates. Terpenoids

was found in all the tested petroleum ether, chloroform and ethanol extract. The presence of sterols was detected in the petroleum ether and chloroform extract.

GC-MS analysis

Spectrums obtained in the GC-MS analysis of all the prepared extracts viz., petroleum ether, chloroform and ethanol extract of the whole plant *I. minor* were matched with the spectra available in the spectral library of NIST version 2.1 to identify the chemical constituents present in the extracts. Peak area, molecular weight and retention time also helped in the identification. Presence of

Table 4: Compounds showed by GC-MS analysis of ethanol extract of *I. minor*.

Sl.No.	RT	Name of the compound	Mol. Formula	Structure	Mol. Wt.	Peak Area %
1	6.54	Butane, 1, 4-bis(methylthio)-	C ₆ H ₁₄ S ₂		150	12.33
2.	9.56	2-Furan carboxaldehyde,5-(hydroxymethyl)-	C ₆ H ₆ O ₃		126	17.50
3.	13.39	Sedoheptulosan	C ₇ H ₁₂ O ₆		196	11.32
4.	16.90	Z,Z-6,2 8-Heptatriactontadien-2-one	C ₃₇ H ₇₀ O		530	3.05
5.	17.15	Trans,cis-1,8 -dimethylspiro[4.5] decane	C ₁₂ H ₂₂		166	1.16
6.	17.34	Cyclohexanol,5-Methyl-2-(1-Methylethyl)-,(1α,2β,5α)-	C ₁₀ H ₂₀ O		156	2.17
7.	19.59	Chloroacetic Acid, Tetradecyl Ester	C ₁₆ H ₃₁ O ₂ Cl		290	2.42
8	22.87	Hentriacontane	C ₃₁ H ₆₄		436	2.039
9.	23.14	1,2-Benzenedicarboxylic Acid, Mono(2-Ethylhexyl) Ester	C ₁₆ H ₂₂ O ₄		278	1.75
10.	25.10	Heneicosane, 3-Methyl	C ₂₂ H ₄₆		310	1.36
11.	25.80	Silane, Trichlorooctadecyl	C ₁₈ H ₃₇ C ₁₃ Si		386	1.50
12.	29.52	Cyclotrisiloxane, Hexamethyl-	C ₆ H ₁₈ O ₃ Si ₃		222	6.06

eight compounds was found in the GC-MS analysis of petroleum ether extract (Figure 2 and Table 2). Presence of six compounds was identified in the chloroform extract (Figure 3 and Table 3) and the ethanol extract showed the presence of twelve compounds (Figure 4 and Table 4).

DISCUSSION

Summarily, result of preliminary qualitative phytochemical evaluation of *I. minor* extracts revealed the presence of various important phytochemicals with significant medicinal values. For example, glycosides are reported to have analgesic and anti-inflammatory, cardio-tonic, anticancer and antimicrobial such as antibacterial, antifungal and antiviral effects. Likewise, the phenolic compounds and tannins are

well known for their antimicrobial, anti-parasitic, antiviral, antioxidant, anti-inflammatory, immunomodulation etc. Terpenoids are renowned for their anti-microbial, anti-allergic, anti-inflammatory, anti-hyperglycemic, anti-spasmodic, and immunomodulatory properties. The previous studies have documented the presence of cardiotoxic, insecticidal, and anti-microbial properties in sterols.

Notably, several of the compounds identified in the GC-MS analysis of the present study have been documented to possess antioxidant, anti-inflammatory, and anticancer properties. For example, Hentriacontane was the major component found in the petroleum ether, chloroform, and ethanol extract of *I. minor* was reported to have anti-inflammatory, antimicrobial, antioxidant




Figure 4: GC-MS examination of ethanol extract of *I. minor*.

and anticancer property.^[19] 2-Furan carboxaldehyde,5-(hydroxymethyl)- present in ethanol extract also documented to have antioxidant and anticancer property. Phytol and 12-oleanen-3-yl acetate, [3 alpha] present in petroleum ether was documented to have anti-inflammatory, antioxidant and anticancer activities.^[20,21]

CONCLUSION

In this study, the whole plant *I. minor* was collected, authenticated, dried and made in to coarse powder. The plant material in powder form was extracted through Soxhlation method, utilizing various solvents such as petroleum ether, chloroform, and ethanol. The dried extract thus obtained was subjected to preliminary phytochemical evaluation and GC-MS investigation. The preliminary phytochemical evaluation, disclosed the absence of alkaloids in all the tested extracts, while the ethanol extract showed the presence of glycosides, phenolic compounds, flavonoids and carbohydrates. Terpenoids was found in all the tested petroleum ether, chloroform and ethanol extract. Petroleum ether and chloroform extract showed the presence of sterols. GC-MS investigation indicated the presence of eight major compounds in the petroleum ether extract, while six major compounds were found in the chloroform extract. Presences of 12 major compounds were found in the ethanol extract of *I. minor*. Summarily, results of the present study revealed the presence of various important phytochemicals with significant medicinal values in the extracts of *I. minor*. This particular GC-MS analysis serves as the initial stage in comprehending the characteristics of active components in this medicinal plant, paving the way for more in-depth research in the future.

ACKNOWLEDGEMENT

We would like to thank our guide Dr. (Sr) Molly Mathew and to Malik Deenar College of Pharmacy, Seethangoli, Kasaragod, Kerala for granting the facilities to carry out this research.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ABBREVIATIONS

GC-MS: Gas Chromatography-Mass Spectrometry; **NIST:** National Institute of Science and Technology.

REFERENCES

- Cragg GM, Newman DJ. Natural product drug discovery in the next millennium. *Pharm Biol.* 2001; 39; Suppl 1:8-17. doi: 10.1076/phbi.39.s1.8.0009, PMID 21554167.
- Suryawanshi HP, Patel MR. Traditional uses, medicinal and phytopharmacological properties of *Erythrina indica* Lam.: an overview. *Int J Res Ayurveda Pharm.* 2011;2(5):1531-3.
- Mandal V, Mohan Y, Hemalatha S. Microwave assisted extraction-An innovative and promising extraction tool for medicinal plant research. *Pharmacogn Rev.* 2007;1(1):7-18.
- Lytovchenko A, Beleggia R, Schauer N, Isaacson T, Leuendorf JE, Hellmann H, et al. Application of GC-MS for the detection of lipophilic compounds in diverse plant tissues. *Plant Methods.* 2009;5:4. doi: 10.1186/1746-4811-5-4, PMID 19393072.
- Janssens SB, Knox EB, Huysmans S, Smets EF, Merckx VS. Rapid radiation of *Impatiens* (*Balsaminaceae*) during Pliocene and Pleistocene: result of a global climate change. *Mol Phylogenet Evol.* 2009;52(3):806-24. doi: 10.1016/j.ympev.2009.04.013, PMID 19398024.
- Yuan YM, Song Y, Geuten K, Rahelivololona E, Wohlhauser S, Fischer E, et al. Phylogeny and biogeography of *Balsaminaceae* inferred from ITS sequences. *Taxon.* 2004;53(2):391-404. doi: 10.2307/4135617.
- Whitfield HL, Toczydlowski RH. Distinguishing *Impatiens capensis* from *Impatiens pallida* (*Balsaminaceae*) using leaf traits. *Botany.* 2020;98(7):361-9. doi: 10.1139/cjb-2020-0022.
- Hatcher PE. *Impatiens noli-tangere* L. *J Ecol.* 2003;91(1):147-67. doi: 10.1046/j.1365-2745.2003.00741.x.
- Delgado Rodríguez FV, Weng Huang NT, Gutierrez AL, Arias Nunez D, Rosales Leiva C. Ethnobotany, pharmacology and major bioactive metabolites from *Impatiens* Genus plants and their related applications. *Pharmacogn Rev.* 2023;17(34):338-81. doi: 10.5530/phrev.2023.17.13.

10. Anwer N, Waqar MA, Iqbal M, Mushtaq M, Sobia A. Phytochemical analysis, free radical scavenging capacity and antimicrobial properties of *Impatiens bicolor* Plant. *Int Food Res J.* 2013;20(1):99-03.
11. Baskar N, Parimala Devi B, Jayakar B. Anticancer studies on ethanol extract of *Impatiens balsamina*. *Int J Res Ayurveda Pharm.* 2012;3(4):631-33.
12. Sebastin V, Gopalakrishnan G, Sreejith M, Anoob Kumar KI. Preliminary phytochemical evaluation and spectral characterization of active constituents in the dried extracts of the whole plant *Argyrea imbricata* (Rox) Sant and Patel. *Asian J Pharm Clin Res.* 2019;12(11):56-60. doi: 10.22159/ajpcr.2019.v12i11.35464.
13. Anoob Kumar KI, Swamivel Manickam M, Sreejith M, Sebastin V. Preliminary phytochemical evaluation, isolation and spectroscopic characterization of constituents in the dried extracts of the whole plant *Crotalaria biflora* (L). *Int J Pharma Bio Sci.* 2020;10(5):124-33. doi: 10.22376/ijpbs/lpr.2020.10.5.P124-133.
14. Arun TS, Rakesh KJ, Jaslin EJ. Preliminary phytochemical and *in vitro* antimicrobial evaluation of *Kedrostis rostrata* plant extracts. *World J Pharm Res.* 2021;7(6):250-55.
15. Kiran KJ, Rakesh KJ, Jaslin EJ. Preliminary studies on phytochemistry and *in vitro* antimicrobial activity of extracts of *Cnidioscolus aconitifolius*. *Eur J Pharm Med Res.* 2021;8(7):424-29.
16. Thasneem CK, Vijayasankar GR, Venkateswarlu BS, Chandira RM, Shanmuganathan S. *In vitro* Antioxidant Activity and Preliminary Phytochemical Evaluation of Different Extracts of *Aerva javanica*. *Pharmacogn Res.* 2022;14(4):468-73. doi: 10.5530/pres.14.4.68.
17. Leslie V, Singh G, Aron S, Harikumar SL. *In vitro* anti-inflammatory evaluation of *Sesbania procumbens* extracts. *Int J Life Sci Pharm Res.* 2022;12(3):70-5. doi: 10.22376/ijpbs/lpr.2022.12.3.P70-75.
18. Gopalakrishnan S, Vadivel E. GC-MS analysis of some bioactive constituents of *Mussaenda frondosa* Linn. *Int J Pharm Biol Sci.* 2011;2(1):313-20.
19. Khajuria V, Gupta S, Sharma N, Kumar A, Lone NA, Khullar M, et al. Anti-inflammatory potential of hentriacontane in LPS stimulated RAW 264.7 cells and mice model. *Biomed Pharmacother.* 2017;92:175-86. doi: 10.1016/j.biopha.2017.05.063, PMID 28549290.
20. Islam MT, Ali ES, Uddin SJ, Shaw S, Islam MA, Ahmed MI, et al. Phytol: a review of biomedical activities. *Food Chem Toxicol.* 2018;121:82-94. doi: 10.1016/j.fct.2018.08.032, PMID 30130593.
21. Abu-Lafi S, Rayan M, Masalha M, Abu-Farich B, Al-Jaas H, Abu-Lafi M, et al. Phytochemical composition and biological activities of Wild *Scolymus maculatus* L. *Medicines (Basel).* 2019;6(2):53. doi: 10.3390/medicines6020053, PMID 31052242.

Cite this article: Sukumaran A, Mathew A, Varghese S, Kannankutty AT. GC-MS Analysis of Bioactive Compounds Present in the Different Extracts of the Whole Plant *Impatiens minor* (DC.) Bennet. *Pharmacogn Res.* 2025;17(1):126-32.