

The heavy metal contents of some selected medicinal plants sampled from different geographical locations

Kofi Annan, Rita A. Dickson, Isaac K. Amponsah, Isaac K. Nooni¹

Department of Pharmacognosy, Faculty of Pharmacy and Pharmaceutical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana, ¹Department of Natural Resource Management, Faculty of Geoinformation Science and Earth Observation, University of Twente, The Netherlands

Submitted: 14-04-2012

Revised: 15-05-2012

Published: 15-04-2013

ABSTRACT

Background: The levels of 5 minerals namely; lead, arsenic, mercury, cadmium, and aluminum were assessed in 10 medicinal plants sampled from 5 different geographical locations to determine the effect of location on the plants' mineral content. **Materials and Methods:** Atomic absorption spectrophotometry (wet digestion) was used for the analyzes, and content of the minerals per sample was expressed as $\mu\text{g/g}$. The levels of minerals were compared to their limit specification for herbs and daily total intake of these minerals. A two-way analysis of variance, which tends to look at the effect of the location and the medicinal plant itself on the plants mineral content, was used in the statistical analysis. **Results:** Lead (Pb) was present in all plant species examined, except *Ocimum gratissimum*. One plant exceeded the maximum safety limit for lead. Cadmium was also detected in some of the medicinal plant species (44%) whilst majority were below the detection limit (0.002) representing 56%. 40% of the plant species exceeded the limit for cadmium. Mercury and arsenic in all the plant species were below the detection limit (0.001). Significant variation existed in mineral content for the various locations ($P \leq 0.05$). **Conclusion:** The findings generally suggest the variation in mineral levels for the various locations. Thus, our study has shown that same species of medicinal plants, growing in different environments, accumulates different levels of heavy metals.

Key words: Geographical locations, heavy metals, medicinal plants, minerals

INTRODUCTION

Herbal medicines have found extensive use in disease treatment, prevention, and management. Due to the immense benefits herbal medicines bring to bear, majority of the world's population in one way or the other depend on them for various health benefits. According to the world health organization report,^[1] there is an estimated 65 to 80% of the world's population relying on traditional (alternative) medicine as their primary form of healthcare. However, the use of herbal medicines has come under scrutiny due to their perceived long term toxicity among other considerations. The causes of the toxicities, which could be attributed to the chemical and mineral contents of various plants, are also linked to the source of the material.^[2]

Address for correspondence:

Dr. Kofi Annan, Department of Pharmacognosy, Faculty of Pharmacy and Pharmaceutical Sciences, Kwame Nkrumah University of science and technology, Kumasi, Ghana.
E-mail: annankofi@yahoo.com

The mineral contents of medicinal plant species used in herbal formulations cannot be over-looked considering the important role these minerals play in the proper functioning of the vital organs as well as in the promotion of the general well-being of the body.^[3] However, they may be toxic if consumed beyond their estimated safe daily intake. Several works have been reported on the phytochemical and biological activities of medicinal plants, although there is little report in regards to the heavy metal contents of these plants. Medicinal herbs can present health risks due to the presence of toxic metals such as Pb, Cd, Al, and Hg, which are hazardous to humans.^[4] The present study was carried out to ascertain the amount of 5 minerals namely; lead (Pb), cadmium (Cd), aluminum (Al), mercury (Hg), and arsenic (As) in 10 common medicinal plants (*Alchornea cordifolia*, *Alstonia boonei*, *Cassia alata*, *Cassia occidentalis*, *Cymbopogon citratus*, *Moringa oleifera*, *Ocimum gratissimum*, *Paullinia pinnata*, *Rauwolfia vomitoria*, and *Taraxacum officinale* leaves) used in the treatment, prevention, and management of diseases and sampled from 5 different geographical locations in Ghana. This may help understand the importance of location in

Access this article online

Website:

www.phcogres.com

DOI: 10.4103/0974-8490.110539

Quick Response Code:



collection and ultimately heavy metal toxicity of medicinal plants. These plants were selected because they are found in most herbal remedies on the Ghanaian market and widely used by individuals and families as they are common in most communities.

MATERIALS AND METHODS

Plant collection

The samples of medicinal plants, which comprised of selected parts of plants commonly used in phytotherapy included; *Alchornia cordifolia* leaves locally known as ‘ogyama,’ *Alstonia boonei* stem bark locally known as ‘onyame dua,’ *Cassia alata* leaves locally known as ‘osempe,’ *Cassia occidentalis* seeds locally known as ‘mmofra brodie,’ *Cymbopogon citratus* leaves locally known as ‘ti-ahaban,’ *Moringa oleifera* leaves, *Ocimum gratissimum* leaves locally known as ‘nunum,’ *Paullinia pinnata* roots locally known as ‘toa-ntini,’ *Rauwolfia vomitoria* stem bark locally known as ‘kakapenpen,’ and *Taraxacum officinale* leaves. The various plant samples were obtained from their specified locations in Kumasi, the Ashanti regional capital in January, 2011 and these were physique garden (PG, faculty of pharmacy and pharmaceutical sciences, Kwame Nkrumah University of Science and Technology (KNUST); located on 6° 40’ 26” N, 1° 33’ 57” W), around KNUST campus (AC; located on 6° 41’ 08” N, 1° 33’ 47” W), by roadside (RS; located on 6° 40’ 11” N, 1° 34’ 22” W), refuse dump (RD; located on 6° 41’ 13” N, 1° 34’ 12” W), and Bobiri forest (FO; located on 6° 43’ 54” N, 1° 25’ 56” W) [Figure 1].

The plant samples were authenticated at the department of herbal medicine, faculty of pharmacy and pharmaceutical sciences, KNUST where voucher specimens are kept [Table 1].

Acid digest and atomic absorption spectrophotometer analysis

An acid digest of each plant species was prepared by oxidizing 0.2 g of the powdered plant sample with nitric/perchloric acid (2:1) mixture (10 ml). One milliliter of the acid digests of each sample was further diluted to 20 ml

due to the corrosive nature of the acid used.^[3] Aliquots of the mixture were used to estimate lead (Pb), mercury (Hg), cadmium (Cd), aluminum (Al), and arsenic (As) by the use of atomic absorption spectrophotometer (model AA240FS). The blank and working standards were first run followed by the samples. Each sample was analyzed twice, and the data reported as a mean of the analyzed samples in µg/g.

Statistical analysis

The data were based on two replicates and subjected to a two-way analysis of variance to bring out the effects of the plants location on the plants mineral content as well the effect the plant itself has on its mineral content. Variations among the locations and plant samples were evaluated by mean of least significance difference (LSD) at 5% level of probability ($P \leq 0.05$). Data analysis was conducted using statgraphics centurion XV I statistical software.

RESULTS

From the study, lead (Pb) was present in all plant species examined with the exception of *O. gratissimum* (AC), which was below the detection limit (<0.010). Cadmium was




Figure 1: Map of KNUST and Bobiri forest showing location of sample points

Table 1: List of plant species under experimentation and their voucher numbers

Species	Family	Common use	Formulation	Voucher number
<i>Alchornia cordifolia</i>	Euphorbiaceae	Septicemia	Decoction	FP1/2011/L038
<i>Alstonia boonei</i>	Apocynaceae	Pains	Decoction	FP1/2011/S003
<i>Cassia alata</i>	Caesalpinaceae	Ringworm	Decoction	FP1/2011/L036
<i>Cassia occidentalis</i>	Caesalpinaceae	Hypertension	Decoction	FP1/2011/SE001
<i>Cymbopogon citratus</i>	Graminae	Fever	Decoction	FP1/2011/L041
<i>Moringa oleifera</i>	Moringaceae	Hypertension	Decoction	FP1/2011/L037
<i>Ocimum gratissimum</i>	Lamiaceae	Flatulence	Decoction	FP1/2011/L040
<i>Paullinia pinnata</i>	Sapindaceae	Impotence	Decoction	FP1/2011/R002
<i>Rauwolfia vomitoria</i>	Apocynaceae	Insomnia	Decoction	FP1/2011/S002
<i>Taraxacum officinale</i>	Compositae	diabetes	Decoction	FP1/2011/L039

also detected in some of the medicinal plant species (44%) whilst majority were below the detection limit (0.002) representing 56%. Aluminum (Al) was found in all of the plant species, except two (4%). Mercury and arsenic in all the plant species were below the detection limit (0.001), representing 100% [Tables 2-5].

DISCUSSION

According to the world health organization (WHO),^[1] about 80% of the world population use plant-based medicines. Most of these plants are collected in the wild, only few are cultivated. A number of mineral elements important to nutrition accumulate in these plants. Other elements such

as Cd, Co, and Pb, which are not used directly by the plant and are detrimental to human health, also accumulate in these plants.^[5,6] The environmental impact of these metals as well as their health effects has been a source of major concern.^[7] Their accumulation in plants highly depends on the availability in the soil.^[7] Heavy metal contamination of medicinal herbal products occurs during cultivation, cross-contamination during processing or their deliberate introduction as therapeutic ingredients. Cultivation in soils containing high concentrations of heavy metals is one mechanism by which heavy metal contamination of herbal products has been documented.^[8]

Medicinal plants have been cited as a potential source of heavy metal toxicity to both man and animals.^[9] The

Table 2: Concentration of heavy metals (µg/g) in *Cassia alata* and *Cassia occidentalis*

Minerals	<i>Cassia alata</i>					<i>Cassia occidentalis</i>				
	RD	AC	RS	FO	PG	RD	AC	RS	FO	PG
Lead	4.450	8.200	17.650	7.450	5.900	4.350	7.850	6.400	7.700	6.000
Cadmium	<0.002	<0.002	0.900	0.150	0.950	<0.002	<0.002	0.350	<0.002	13.700
Aluminum	65.900	100.45	8.950	14.250	106.3	67.150	98.750	4.800	20.650	96.275
Mercury	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Arsenic	<0.001	0.001	0.002	<0.001	<0.001	<0.001	0.001	<0.001	0.001	<0.001

RD=Refuse dump; AC=Around campus; RS=Roadside; FO=Forest; PG=Physique garden values with<=below detection limit of the spectrophotometer

Table 3: Concentration of heavy metals (µg/g) in *Rauwolfia vomitoria* and *Paulinia pinnata*

Minerals	<i>Rauwolfia vomitoria</i>					<i>Paulinia pinnata</i>				
	RD	AC	RS	FO	PG	RD	AC	RS	FO	PG
Lead	1.550	9.250	7.750	4.600	4.750	2.700	8.450	4.950	9.050	6.700
Cadmium	<0.002	0.850	<0.002	<0.002	<0.002	<0.002	0.650	<0.002	<0.002	<0.002
Aluminum	82.650	101.800	23.350	23.200	105.525	76.650	104.25	23.800	15.850	12.400
Mercury	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Arsenic	<0.001	0.001	0.001	<0.001	<0.001	<0.001	0.001	<0.001	0.001	<0.001

RD=Refuse dump; AC=Around Campus; RS=Roadside; FO=Forest; PG=Physique Garden values with<=below detection limit of the spectrophotometer

Table 4: Concentration of heavy metals (µg/g) in *Moringa oleifera* and *Taraxacum officinale*

Minerals	<i>Moringa oleifera</i>					<i>Taraxacum officinale</i>				
	RD	AC	RS	FO	PG	RD	AC	RS	FO	PG
Lead	5.350	3.250	3.400	6.400	3.900	6.250	1.650	4.650	7.250	4.800
Cadmium	0.550	0.500	0.650	0.350	<0.002	14.050	<0.002	1.500	<0.002	0.950
Aluminum	77.150	110.475	33.850	20.750	16.450	94.375	105.85	28.400	11.500	21.450
Mercury	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Arsenic	0.001	<0.001	<0.001	<0.001	<0.001	0.002	<0.001	<0.001	<0.001	<0.001

RD=Refuse dump; AC=Around campus; RS=Roadside; FO=Forest; PG=Physique garden values with<=below detection limit of the spectrophotometer

Table 5: Concentration of heavy metals (µg/g) in *Ocimum gratissimum* and *Alstonia boonei*

Minerals	<i>Ocimum gratissimum</i>					<i>Alstonia boonei</i>				
	RD	AC	RS	FO	PG	RD	AC	RS	FO	PG
Lead	4.600	<0.010	6.650	3.250	5.550	6.400	4.950	8.350	4.900	8.650
Cadmium	<0.002	<0.002	1.800	<0.002	2.250	<0.002	<0.002	1.450	<0.002	1.800
Aluminum	94.275	107.60	25.950	31.650	3.550	94.425	105.800	24.700	36.000	<0.030
Mercury	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Arsenic	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.001	<0.001	0.001

RD=Refuse dump; AC=Around campus; RS=Roadside; FO=Forest; PG=Physique garden values with<=below detection limit of the spectrophotometer

most common heavy metals implicated in human toxicity include lead, mercury, arsenic, and cadmium, although aluminum and cobalt may also cause toxicity. Therefore, the world health organization recommends that medicinal plants, which form the raw materials for most herbal remedies, should be checked for the presence of heavy metals. However, majority of people, living in areas where these plants grow, harvest them locally for personal or family use without checking for heavy metal accumulation. The general notion that medicinal plants are safe and devoid of heavy metal toxicity could be misconstrued. From the study, the levels of these metals differed in the same plant collected from different geographical locations. For example, the levels of lead in *Cassia alata* varied from 17.7 to 4.45 µg/g for the 5 collection sites [Table 2]. That for *Cassia occidentalis* and *Rauwolfia vomitoria* varied between 7.85-4.35 and 9.25-1.55 µg/g, respectively (Table 2 and 3). Similarly, that of aluminum varied between 105.53-23.3 for *Rauwolfia vomitoria* and 104.25-12.4 µg/g for *Paullinia pinnata* [Table 3]. The levels of heavy metals also varied for different plants collected from the same location. Uptake of metals by plants is influenced by a number of factors including metal concentrations in soils, cation exchange capacity, soil pH, organic matter content, types and varieties of plants, and plant age. However, the prevailing factor is the concentration of the metal in the soil^[10] and thus the existing environmental conditions.

Lead is considered very harmful for plants, animals, and particularly for microorganisms.^[7] It has no physiologic role. The highest level of lead occurred in *Cassia alata* harvested from the roadside (17.65 µg/g) [Figure 2]. This exceeded the WHO standard of 10 µg/g for lead in raw materials for herbal medicines.^[1] This could possibly be due to its accumulation in the soil from vehicular exhaust fumes. In a large number of the samples, the level of lead was just below the allowable limit, suggesting that toxicity could arise from long term use of these plants. Levels of lead beyond the permissible values or long term use of these plants could lead to toxicity characterized by colic, anemia, headache, convulsions and chronic nephritis of the kidneys, brain damage, and central nervous system disorders.^[7,11]

The levels of mercury and arsenic, in all the medicinal plant samples in different geographical locations, were found to be below the minimum detectable limit (<0.001 µg/g) [Table 2 and 6]. All values were below the permissible values of 5 and 0.2 µg/g for arsenic and mercury, respectively.^[1] Thus, human activities in the areas, where these plants were collected, do not lead to the accumulation of arsenic and mercury in the soils. Mercury levels in humans above the allowable values have been associated with male infertility, inhibition of endogenous antioxidant enzymes, and brain damage among others.^[12] Supraoptimal levels of arsenic are often fatal. Other toxic symptoms include peripheral polyneuropathy, liver cirrhosis, visual disturbance, and blindness.^[13] Similarly, aluminum levels in all plant samples from the same area in different geographical locations varied from 4.8 to 108 µg/g [Tables 2-6]. All values were within allowable limits for aluminum, which is not used in any biochemical process in living systems. It accumulates in the brain and, to a lesser extent, in the bones.^[14] The levels of aluminum in medicinal plants in different geographical locations must be checked as long term use could lead to toxicity.

The level of cadmium, in all 10 plant samples, exceeded the WHO recommended value of 0.3 µg/g for some geographical locations, whereas for other collection sites, the levels were below the minimum detectable values (<0.001 µg/g) [Figure 3]. The levels of cadmium from plant samples collected from the roadside exceeded the maximum safety limit, indicated by the red line in Figure 3, possibly due to contamination from vehicular activities. Also, most medicinal plants collected around the university campus and the physic garden accumulated cadmium above the recommended levels in medicinal plants [Figure 3] with the highest occurring in *C. occidentalis* (13.7 µg/g). About 70% of medicinal plants harvested from the physic garden (within KNUST campus) exceeded the maximum safety limit for cadmium. The most common sources for cadmium in soil and plants are phosphate fertilizers, sewage sludge application, and combustion of fossil fuels. Thus, there is the need for a proper and efficient waste disposal system in and around the university community as cadmium levels above permissible values may result in irreversible kidney damage.^[15]

Table 6: Concentration of heavy metals (µg/g) in *Cymbopogon citratus* and *Alchornea cordifolia*

Minerals	Cymbopogon citratus					Alchornea cordifolia				
	RD	AC	RS	FO	PG	RD	AC	RS	FO	PG
Lead	8.050	2.250	6.500	5.400	7.950	6.350	5.450	7.400	6.950	5.450
Cadmium	<0.002	<0.002	1.200	<0.002	1.250	<0.002	<0.002	0.700	<0.002	0.700
Aluminum	89.575	108.075	44.400	26.350	<0.030	100.275	97.150	38.850	15.950	10.300
Mercury	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Arsenic	0.001	<0.001	<0.001	<0.001	0.001	<0.001	<0.001	<0.001	0.001	<0.001

RD=Refuse dump; AC=Around campus; RS=Roadside; FO=Forest; PG=Physique garden values with <=below detection limit of the spectrophotometer




Figure 2: Levels of lead in plant samples in different geographical locations. RD = Refuse dump, AC = Around campus, RS = Roadside, FO = Forest, PG = Physique garden values with <=below detection limit of the spectrophotometer




Figure 3: Levels of cadmium in plant samples in different geographical locations. RD=Refuse dump, AC=Around xampus, RS=Roadside, FO=Forest, PG=Physique garden values with <=below detection limit of the spectrophotomete

CONCLUSION

Our study has shown that same species of medicinal plants, growing in different geographical locations, cumulates different levels of heavy metals. Concentration of heavy metals also differed for different plant species collected from the same geographical location. Amounts of heavy metals detected in plants collected from the Bobiri forest reserve were within permissible limits, whereas the levels

in other geographical locations exceeded the recommended values. Thus, medicinal plants for the formulation of herbal remedies should be harvested from pollution-free natural habitat. Our findings further indicate that the medicinal plants, used for local or pharmaceutical purposes, should be collected from areas not contaminated with heavy metals. It is, therefore, advised that the metal content in medicinal plants be checked for levels of heavy metals before their use for local and pharmaceutical purposes.

ACKNOWLEDGEMENT

The authors are grateful to the technical staff of Atomic Energy Commission, Accra, Ghana for their assistance in this work.

REFERENCES

1. WHO. Quality control methods for medicinal plant materials. WHO Geneva Switzerland. 1998, available at <http://whqlibdoc.who.int/publications/1998/9241545100.pdf>.
2. Lekouch N, Sedki A, Nejmeddine A, Gamon S. Lead and traditional Moroccan pharmacopoeia. *Sci Total Environ* 2001;280:39-43.
3. Annan K, Kojo AI, Cindy A, Samuel AN, Tunkumgnen BM. Profile of heavy metals in some medicinal plants from Ghana commonly used as components of herbal formulations. *Pharmacognosy Res* 2010;2:41-4.
4. Yap C K, Fitri M, Mazyhar Y and Tan SG. Effects of Metal-contaminated Soils on the accumulation of heavy metals in different parts of *Centella asiatica*: A Laboratory Study. *Sains Malaysiana* 2010;39:347-52.
5. Baker AJM and Brooks RR. Terrestrial higher plants, which hyperaccumulate metallic elements—A review of their distribution, ecology and phytochemistry. *Biorecovery* 1989;1:81-126.
6. Lasisi AA, Yusuff AA, Ejelonu BC, Nwosu EO, Olayiwola MA. Heavy metals and macronutrients content in selected herbal plants of Nigeria. *International Jour Chern* 2005;15:147-54.
7. Khan MA, Ahmad I and Rahman I. Effect of environmental pollution on heavy metals content of *Withania somnifera*. *Journal of the Chinese Chemical Society* 2007;54:339-43.
8. Quig D. Cysteine metabolism and metal toxicity. *Altern Med Rev* 1998;3:262-70.
9. Dwivedi SK, Dey S. Medicinal herbs: A potential source of toxic metal exposure for man and animals in India. *Arch Environ Health* 2002;57:229-31.
10. Jun MC. Heavy metal concentrations in soils and factors affecting metal uptake by plants in the vicinity of a Korean Cu-W Mine. *Sensors* 2008;8:2413-23.
11. Zheljzkov VD. Heavy metal content in some essential oils and plant extracts. Amhers, USA: International Society for Horticultural Science; 1995. p. 429.
12. Choy CM, Lam CW, Cheung LT, Briton-Jones CM, Cheung LP, Haines CJ. Infertility, blood mercury concentrations and dietary seafood consumption: A case-control study. *BJOG* 2002;109:1121-5.
13. Ling LJ, Clark RF, Erickson TB, Trestrail JH. *Toxicology secrets*. Philadelphia: Hanley and Belfus Inc; 2001. p. 153-69.
14. Cassel I. Aluminum Toxicity: A misdiagnosed epidemic (Part 1). *Idaho Observer*; 2007.
15. Saied S, Zahir H. and Siddique A. Heavy metal levels in commonly used traditional medicinal plants. *J chem Soc Pak* 2010;32:737-43.

Cite this article as: Annan K, Dickson RA, Amponsah IK, Noonik IK. The heavy metal contents of some selected medicinal plants sampled from different geographical locations. *Phcog Res* 2013;5:103-8.

Source of Support: Nil, **Conflict of Interest:** None declared.