













- fruits against intestinal bacteria. *J Food Prot* 2006;69:2205-9.
3. Vonshak A, Barazani O, Sathiyamoorthy P, Shalev R, Vardy D, Golan-Goldhirsh A. Screening South Indian medicinal plants for antifungal activity against cutaneous pathogens. *Phytother Res* 2003;17:1123-5.
  4. Shukla YN, Srivastava A, Kumar S, Kumar S. Phytotoxic and antimicrobial constituents of *Argyrea speciosa* and *oenothera biennis*. *J Ethnopharmacol* 1999;67:241-5.
  5. Koga T, Moro K, Nakamori K, Yamakoshi J, Hosoyama H, Kataoka S, *et al*. Increase of antioxidative potential of rat plasma by oral administration of proanthocyanidin-rich extract from grape seeds. *J Agric Food Chem* 1999;47:1892-7.
  6. Burns J, Gardner PT, O'Neil J, Crawford S, Morecroft I, McPhail DB, *et al*. Relationship among antioxidant activity, vasodilation capacity, and phenolic content of red wines. *J Agric Food Chem* 2000;48:220-30.
  7. Vandendool H, Kratz PD. A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *J Chromatogr* 1963;11:463-71.
  8. Harborne JC. *Phytochemical methods: A guide to modern techniques of plant analysis*. 2<sup>nd</sup> ed. London, New York: Chapman and Hall Ltd.; 1984.
  9. Andrews JM. BSAC standardized disc susceptibility testing method. *J Antimicrob Chemother* 2001;4:43-57.
  10. Barsi DF, Fan SH. The potential of aqueous and acetone extracts of galls of *Quercus infectoria* as antibacterial agents. *Indian J Pharmacol* 2005;37:26-9.
  11. Juang LJ, Sheu SJ, Lin TC. Determination of hydrolyzable tannins in the fruit of *Terminalia chebula* Retz. by high-performance liquid chromatography and capillary electrophoresis. *J Sep Sci* 2004;27:718-24.
  12. Lin-Jeng J, Shuenn-Jyi S, Ta-Chen L. Determination of hydrolysable tannins in the fruit of *Terminalia chebula* Retz. by high-performance liquid chromatography and capillary electrophoresis. *J Sep Sci* 2004;27:718-24.
  13. Sangeetha J, Vijayalakshmi K. Determination of bioactive components of ethyl acetate fraction of *Punica granatum* rind extract. *Int J Pharm Sci Drug Res* 2011;3:116-22.
  14. Moridani MY, Scobie H, O'Brien PJ. Catechin metabolism: Glutathione conjugate formation catalysed by tyrosinase, peroxidase, and cytochrome 450. *Chem Res Toxicol* 2001;14:841-8.
  15. Skaper SD, Fabris M, Ferrari V, Daile Carbonare M, Leon A. Quercetin protects cutaneous tissue-associated cell type including sensory neurons from oxidative stress induced by glutathione depletion: Cooperative effects of ascorbic acid. *Free Radic Biol Med* 1997;22:669-78.
  16. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationship of flavonoids and phenolic acid. *Free Radic Biol Med* 1996;20:933-56.
  17. Chen ZP, Schell JB, Ho CT, Chen KY. Green tea epigallocatechin gallate show a pronounced growth inhibitory effect on cancerous cells but not on their normal counterparts. *Cancer Lett* 1998;1129:173-9.
  18. Ultee A, Bennik MH, Moezelaar R. The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Appl Environ Microbiol* 2002;68:1561-8.
  19. Brown WH. *Introduction to organic chemistry*. 3<sup>rd</sup> ed. Boston, Mass: Willard Grant Press; 1975.
  20. Shelef LA. Antimicrobial effects of spices. *J Food Saf* 1983;6:29-44.
  21. Kocacaliskan I, Talan I, Terzi I. Antimicrobial activity of catechol and pyrogallol as allelochemicals. *Z Naturforsch C* 2006;61:639-42.
  22. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999;12:564-82.

**Cite this article as:** Singh G, Kumar P. Extraction, gas chromatography-mass spectrometry analysis and screening of fruits of *Terminalia chebula* Retz. for its antimicrobial potential. *Phcog Res* 2013;5:162-8.

**Source of Support:** UGC, **Conflict of Interest:** Department of Botany, University of Rajasthan.