

Evaluation of Antioxidant and Hepatoprotective Activity of Fruit Rind Extract of *Garcinia dulcis* (Roxburgh) Kurz

Nabajyoti Gogoi, Ankur Gogoi, Bijoy Neog¹, Dibyojyoti Baruah¹, Khumanthem Deepak Singh²

Centre for Biotechnology and Bioinformatics Dibrugarh University, ¹Department of Life Sciences, Plant Molecular Biology Laboratory, Dibrugarh University, ²Department of Pharmaceutical Science, Dibrugarh University, Dibrugarh, Assam, India

ABSTRACT

Background: *Garcinia* spp. belongs to the family *Clusiaceae* has been traditionally used for the treatment of many ailments including the liver damage. *Garcinia dulcis* found in North Eastern region of Assam; India can be a potential candidature to combat different ailments. Objective: The present work has been designed in such a way to appraisal the antioxidant and hepatoprotective activity of fruit rind extract of this plant. **Materials and Methods:** The antioxidant activity was investigated through the various *in vitro* models, namely, 2,2-diphenyl-1-picrylhydrazine, 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid, nitrite oxide. Phytochemical investigation for total phenolic and flavonoids contents were carried out by standard protocol. For the evaluation of hepatoprotective activity, albino Wistar rats were divided into five groups, five animals per group and activity was determined by measuring the contents of liver function marker enzymes such as serum glutamate oxaloacetate transaminase (SGOT), serum glutamate pyruvate transaminase (SGPT), serum alkaline phosphatase (ALP), and biochemical parameter, that is, Bilirubin and total protein. Histopathology observation of liver sections was conducted. **Results:** Phytochemical investigation revealed the presence of both phenolic and flavonoid groups in the extract in a significant amount. Antioxidant activity of the plant extract was observed in all models and percentage of inhibition was dose-dependent. Intoxicated with carbon tetrachloride, elevated the liver function enzymes, bilirubin, and suppressed the production of total protein. Pretreatment with the extract decreased the SGOT, SGPT, ALP, and bilirubin level significantly and increased the production level of total protein in a dose-dependent manner. The histopathological observation supported the hepatoprotective potentiality of the extract. **Conclusion:** The results indicate that fruit rind part of *G. dulcis* is nontoxic and the plant can utilize as an antioxidant source. The plant has a protective agent for liver damages and other diseases caused by free radicals.


Key words: Antioxidant, free radicals, *Garcinia dulcis*, hepatoprotective

SUMMARY

- *In vitro* antioxidant and *in vivo* hepatoprotective activity was evaluated
- Methanolic extract was subjected to quantify the both phenolic and flavonoid contents. The extract showed the significant amount of both phenolic and flavonoids contents. The extract showed the free radical scavenging activity in 2,2-diphenyl-1-picrylhydrazine, 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid and nitrite oxide models
- The extract was administrated against the carbon tetrachloride intoxicated animal models to evaluate the hepatoprotective activity by determining the level of liver marker enzymes such as serum glutamate pyruvate transaminase,

SGOT, alkaline phosphatase and biochemical parameter such as protein and bilirubin. Pretreatment with the extract reversed the elevated level of the enzymes and increased the protein level in a dose-dependent manner

- The histopathological observations of the liver sections supported the hepatoprotective activity of the extract
- The present study revealed that the *Garcinia dulcis* extract is a good candidature for preventing liver damage and other disease caused by free radicals.



Abbreviations Used: DPPH: 2,2-diphenyl-1-picrylhydrazine, ABTS: 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid, SGOT: Serum glutamate oxaloacetate transaminase, SGPT: Serum glutamate pyruvate transaminase, ALP: Serum alkaline phosphatase.

Correspondence:

Prof. Bijoy Neog,
Department of Life Sciences, Plant Molecular Biology Laboratory, Dibrugarh University, Dibrugarh, Assam, India.
E-mail: bijoyneog@dibru.ac.in
DOI: 10.4103/0974-8490.210330

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Free radicals are produced in the human bodies which cause different disorders such as liver injury, Alzheimer's diseases, atherosclerosis, cardiovascular diseases, diabetes, and rheumatoid arthritis. However, human body developed their own defense mechanism to remove or neutralized the free radicals with the help of antioxidant enzymes such as catalase, superoxide dismutase, and glutathione peroxidase.^[1-3] However, due to heavy exposure to external environmental pollutants, such as smoke, ultraviolet (UV) radiation, and pesticides, the self-defense mechanism is sometimes insufficient to remove the excessive free radicals. Therefore, supplementation of phytochemicals

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Gogoi N, Gogoi A, Neog B, Baruah D, Singh KD. Evaluation of antioxidant and hepatoprotective activity of fruit rind extract of *Garcinia dulcis* (Roxburgh) Kurz. *Phcog Res* 2017;9:266-72.

with antioxidant properties has become a considerable attention at present scenario.^[4]

The liver is the largest and most vital organ of the human body and involved in almost all the biochemical pathways such as in secretion, growth, supply of nutrient, energy, and reproduction. The liver is the primary site of intense metabolism and excretion. The major function of liver is the metabolism of carbohydrate, fats and protein; and storage of energy for future use. Therefore, protection of liver from the toxic products is a serious concern for well-being. However, due to consumption of alcohol, exposure of environmental toxin, over drugs consumption lead the liver in various ailments like hepatitis, cirrhosis, etc.^[5,6] Carbon tetrachloride (CCl₄) is a widely used hepatotoxic chemical to induce the liver and tissue damage in laboratory animal experiment.^[7,8] The main principle of CCl₄ induced liver damage is the lipid peroxidation and decreases the antioxidant enzyme system by producing free radicals.^[9] Many synthetic drugs used in the treatment of liver disorder are inadequate and sometimes create critical side effect.^[10] Therefore, nowadays, there is growing interest in evaluation of hepatoprotective activity of traditionally used herbal medicine in a scientific manner and their treatment is safe because they are natural.^[11]

Garcinia Linn. belongs to the family of *Clusiaceae*, (Thekera, in Assamese, India) is traditionally used for cure of several diseases such as dysentery, liver damage, after childbirth complex and fever.^[12] The pericarp extract of this family is rich in the both phenolic and flavonoids contents and plays a significant role in the scavenging of free radicals in the different *in vitro* antioxidant models.^[13] *Garcinia dulcis* (Tepor tenga, in Assamese, India) was reported from Dibrugarh, Assam, India by Begum *et al.*^[14] which contains considerable medicinal properties, as per traditional knowledge system.

Therefore, the present study was undertaken to evaluate the *in vitro* antioxidant potential and hepatoprotective activity of fruit rind extract of *G. dulcis* against the CCl₄ induced albino rats.

MATERIALS AND METHODS

Plant material collection and preparation of extract

Fresh fruits were collected from Jokai Botanical Garden, Dibrugarh, Assam, India, in the month of September to November 2014 and identified by Department of Life Sciences, Dibrugarh University. A reference specimen has been retained in the herbaria in the Department of Life Sciences for future reference.

The fruit rind was dried in hot air oven by maintaining the temperature between 25°C and 35°C. The oven-dried sample was then powdered and 50 g of the sample immersed in 100 ml of 80% methanol in a shaking flask condition for 24 h (at 25°C) and later filtered through Whatman No. 1 filter paper. The extract was then concentrated with the help of the rotary evaporator under pressure, and finally, extract was stored in the refrigerator for further experiment.

Chemicals

All the chemicals and solvents were of analytical grade and procured from the Merck Millipore. 2,2-diphenyl-1-picrylhydrazine (DPPH), ascorbic acid, 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS), naphthyl ethylenediamine dihydrochloride were collected from the Sigma-Aldrich (Germany). Other chemicals such as Silymarin, CCl₄ were from Merck Millipore. Standard kits for SGOT, serum glutamate pyruvate transaminase (SGPT), serum alkaline phosphate (SALP), and total bilirubin were purchased from the Span Diagnostics Ltd., India.

Phytochemical investigation

Plant extract was subjected toward the quantification of total phenolic and flavonoids contents. Gallic acid and Rutin were considered standard

compounds, respectively and the result was expressed in standard equilibrium.

Estimation of total phenolic content

Half milliliter of each plant extract (1000 µg/ml) was mixed with the 2.5 ml of Folin-Ciocalteu reagent (diluted 1:10 ratio with distilled water) and 1.5 ml of sodium carbonate (20% w/v). The mixtures were shaken thoroughly using vortex and made up to the volume of 10 ml with double distilled water. The mixtures were then kept for 90 min. The absorbance was measured at 760 nm using UV - Spectrophotometer. Different concentration (10–50 µg/ml) of Gallic acid was used for the preparation of the standard curve. The total phenol content was expressed in Gallic acid equivalent in mg/g of plant extract.^[15]

Estimation of total flavonoids content

Half milliliter of each plant extract (1000 µg/ml) was mixed with 1.5 ml of methanol (75% v/v), 0.1 ml (1 M) of potassium acetate and keep for 1 min. After 1 min incubation, 0.1 ml (10% w/v) aluminum chloride was added and made up the volume 3 ml with double distilled water and kept the mixture for 30 min in dry and shady place. The absorbance was measured at 415 nm. Rutin was used as standard flavonoids for the preparation of calibration curve and total flavonoids was expressed as rutin equivalent in mg/g of plant extract.^[16]

Evaluation of antioxidant activity

Antioxidant activity was evaluated in different *in vitro* assay models.

2,2-diphenyl-1-picrylhydrazine radical scavenging assay

DPPH is a stable free radical which is usually converted into 1, 1 dihydroxyl 2-picryl hydration. Whenever it reacts with antioxidant, the degree of conversion can be measured by spectrophotometric method. A volume of 2 ml of different concentration (100, 500, 1000 µg/ml) of the extract and different concentration (100, 500, 1000 µg/ml) of standard solution (Ascorbic acid) were added to the freshly prepared 2 ml of DPPH solution (0.2 mM). After incubation of 30 min in the dark place, absorbance was taken in 517 nm using UV-spectrophotometer.^[17] The experiment was performed in triplicate and percentage of inhibition was calculated by the following formula.

$$\text{Percentage of inhibition (\%)} = \frac{\text{Absorbance of control} - \text{Absorbance of test sample}}{\text{Absorbance of control}} \times 100\%$$

2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid radical scavenging assay

The ABTS radical scavenging assay was carried out by the method of Re *et al.*^[18] with minor modification. The ABTS radical was formed by the oxidation of ABTS with potassium persulfate. The ABTS cation solution was prepared by mixing of 5 ml ABTS (7 mm) with 100 µl of potassium persulfate and incubated for 12 h. For the preparation of working solution, a necessary amount of the previous solution was added to the phosphate buffer saline (PBS) solution till the absorbance was 0.70 ± 0.001 at 734 nm. The freshly prepared working solution was then kept in the dark for 30 min. One milliliter of different concentrations of plant extract was mixed with 1 ml of working solution and mixed thoroughly; kept in the dark for 10 min. The absorbance was measured at 734 nm. The percentages of inhibition were calculated by above-mentioned formula.

Nitrite oxide radical inhibition assay

Nitric oxide was generated by dissolving the sodium nitroprusside in an aqueous solution which was measured by Griess reaction. Scavengers of nitric oxide compete with oxygen to help reduce the production of nitride, oxide.^[19] The reaction (6 ml) mixer was prepared by mixing of sodium nitroprusside (10 mM, 4 ml), 1 ml of different concentration of sample or standard (dissolved in dimethyl sulfoxide) and PBS (1 ml PH 7.4) and kept the mixer for reaction at 25°C for 150 min. After the reaction, 0.5 ml of reaction mixer was removed and transferred to the Griess reagent and incubated for 30 min at diffused light. Griess reagent was prepared by mixing of sulfanilic acid (1 ml, 0.33% w/v) and naphthylethylenediamine dihydrochloride (1 ml, 0.1% w/v). The absorbance was measured at 540 nm.^[20] The percentage of inhibition was calculated above mention formula.

In vivo animal experiment

Experimental animals

Wistar albino female rats, weighting 80–120 g were used for hepatoprotective experiment. Animals were maintained under standard condition (12 h light/dark cycle; 25°C), fed with standard feed and water *ad libitum*, at the Central Animal House of Dibrugarh University. Animals were acclimatized to the animal house conditions for 15 days before conducting the experiment. The experimental protocol was reviewed and approved by Dibrugarh University animal ethical committee (approved no. IAEC/DU/98, Regd. no. 1576/GO/a/11/CPCSEA) and animals were cared and handled with before the regulation of CPCSEA.

Acute toxicity study

Acute toxicity of methanol fruit rind extract was of *G. dulcis* was conducted on Albino rats, according to the OECD guideline No. 425.^[21] Animals were fasted for the 12 h, and the extract of 2000 mg/kg (single dose) was administrated orally, and observed for initially for 6 h and conducted the experiment for 14 days to understand mortality, general behavior changes, discomforts and nervous manifestations.

Carbon tetrachloride induced hepatoprotective activity

Albino rats were divided randomly into the five groups and containing six animals. The experiment was conducted for the 7 days.

Group I: Positive control (normal) group was administrated with normal saline, 0.9% NaCl (5 ml/kg body weight, po), for 7 days.

Group II: Negative controlled CCl₄ induced group, was administrated with a single dose of normal saline daily, 0.9% of NaCl (5 ml/kg, po) and CCl₄/olive oil (1:1 v/v, 1 ml/kg, ip) on the alternate days for 7 days.

Group III: Standard group, was administrated with a single daily dose of Silymarin (50 mg/kg, po) and CCl₄/olive oil (1:1 v/v, 1 ml/kg, ip) on every alternate days for 7 days.

Group IV: Low-dose test group, was administrated with methanolic extract (200 mg/kg, po) single dose daily and CCl₄/olive oil (1:1 v/v, 1 ml/kg, ip) in the alternate days of 7 days.

Group V: High-dose test group, was administrated with methanolic extract (400 mg/kg, po) single dose daily and CCl₄/olive oil (1:1 v/v, 1 ml/kg, ip) in the alternate days for 7 days.^[22]

Assessment of hepatoprotective activity

Biochemical estimation

On the 8 days, after 12 h administration of the last dose, bloods were collected from the retro-orbital plexus using ether as anesthesia. The animals were then sacrificed by cervical decapitation. Blood was also withdrawn by intracardiac puncture. Serum was separated from the

fresh blood by centrifugation at 5000 rpm for 10 min and stored at 4°C until use. The conventional liver biochemical test from the blood serum, namely, Serum Glutamate Pyruvate Oxaloacetate Transaminase (SGOT), SGPT were estimated by the methods of Reitman and Frankel^[23] and SALP was estimated by the methods of King and Kind^[24] using standard kit from Span Diagnostics Ltd., India. Total protein and bilirubin contents were estimated by Lowry *et al.* and Malloy and Evelyn methods, respectively.^[25,26]

Histopathological studies

After collection of blood by intracardiac puncture, liver was excised from the different group, washed with normal saline and dried with blotting paper. It was fixed with 10% formalin. The fixed liver was then dehydrated in graded alcohol (30%–100%) and embedded in paraffin. Microtome sections (0.5 μ) were prepared and stained with hematoxylin-eosin dye and finally examined under microscope (×40) for histopathological changes for the hepatoprotective assessment.^[27]

Statistical analysis

The data were expressed in mean ± standard error, (*n* = 3, 6). All data were analyzed by one-way ANOVA followed by Tukey test. The graphical representations were prepared using Sigma Plot software (Systat software Inc. version 13, San Jose, USA).

RESULTS

Phytochemical investigation

Total phenolic and flavonoids contents

The quantification of total phenolic and flavonoids contents of *G. dulcis* is summarized in Table 1. The results were expressed in Gallic acid equivalent (mg/g of extract) and Rutin equivalent (mg/g extract), respectively.

Antioxidant activity investigation

2,2-diphenyl-1-picrylhydrazine radical scavenging assay

The inhibition percentage of DPPH radical scavenging activity of three concentrations of the rind extract is presented in Figure 1. The rind extract exhibited dose-dependent inhibition on DPPH models. The inhibition of DPPH free radicals was increased with increasing the concentration of the plant extract. The maximum inhibition of the extract was 56.44% ± 3.45%, while ascorbic acid was 231.86% ± 2.18% at 1000 μg/ml.

2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid radical scavenging assay

The inhibition percentage ABTS radical scavenging of the three concentrations of plant extract is presented in Figure 2. As illustrated in Figure 2, the scavenging activity of the extract was dose dependent and increased the percentage of inhibition with increasing the concentrations. The high inhibition was found at 1000 μg/ml concentration, i.e.: 64.09% ± 1.21%, while it was found 123.1% ± 2.10% in the case of Ascorbic acid.

Table 1: Estimate value of total phenolic and flavonoids contents of *Garcinia dulcis*

Plants extract	Total phenolic content (GAE mg/g dry extract)	Total flavonoids content (RE mg/g dry extract)
<i>Garcinia dulcis</i>	56.52±2.94	84.88±3.91

Values are mean±SEM of three replicates. GAE: Gallic acid equivalent; RE: Rutin equivalent; SEM: Standard error mean

Nitrite oxide radical inhibition assay

Nitride oxide radicals inhibition of the rind extract was also found to be dose-dependent. The inhibition was increased with increasing of concentrations of the plant extract. The maximum inhibition was recorded $56.15\% \pm 0.972\%$, while in ascorbic acid, it was 418.99 ± 5.35 , at $1000 \mu\text{g/ml}$. The result is presented in Figure 3.

In vivo animal experiment

Acute toxicity

There is no mortality among the animals and recorded no toxicity, and abnormal behavior observed at the dose level 2000 mg/kg . The results

suggest that the rind extract is safe at the dose of 2000 mg/kg body weight. Therefore, administrated dose was selected at $200, 400 \text{ mg/kg}$, po for further experiments.

Effect of rind extract on serum marker enzyme, total protein, and bilirubin

The serum marker SGOT, SGPT, SALP, and bilirubin were significantly elevated in the content of CCl_4 intoxicated animals. However, the total protein level was reduced in CCl_4 intoxicated animals. In contrast, pretreatment with rind extract ($200, 400 \text{ mg/kg}$) and Silymarin (50 mg/kg) exhibited a reversed parameter level significantly in a dose-dependent manner. The results are presented in Figures 4-8.




Figure 1: The effect of *Garcinia dulcis* fruit rind extract and known antioxidant (ascorbic acid) in 2,2-diphenyl-1-picrylhydrazine model. Values are mean \pm standard error of three replicates




Figure 2: The effect of *Garcinia dulcis* fruit rind extract and known antioxidant (ascorbic acid) in 2,2'-azino-bis-3-ethylbenzothiazoline-6-sulfonic acid model. Values are mean \pm standard error of three replicates




Figure 3: The effect of *Garcinia dulcis* fruit rind extract and known antioxidant (ascorbic acid) in nitrogen oxide model. Values are mean \pm standard error of three replicates




Figure 4: Effect of *Garcinia dulcis* extract on SGOT enzyme level in experimental rat. ^a $P < 0.001$ when negative group was compared with normal group; ^x $P < 0.001$, ^y $P < 0.001$, ^z $P < 0.001$ when test groups were compared with negative group




Figure 5: Effect of *Garcinia dulcis* extract on serum glutamate pyruvate transaminase enzyme level in experimental rats. ^a*P* < 0.001 when negative group was compared with normal group; ^x*P* < 0.001, ^y*P* < 0.001, ^z*P* < 0.001 when test groups were compared with negative group




Figure 6: Effect of *Garcinia dulcis* extract on alkaline phosphatase enzyme level in experimental rats. ^a*P* < 0.001 when negative group was compared with normal group; ^x*P* < 0.001, ^y*P* < 0.001, ^z*P* < 0.001 when test groups were compared with negative group




Figure 7: Effect of *Garcinia dulcis* extract on total protein level in experimental rats. ^a*P* < 0.001 when negative group was compared with normal group; ^x*P* < 0.001, ^b*P* < 0.05 when test groups were compared with negative group




Figure 8: Effect of *Garcinia dulcis* extract on total bilirubin level in experimental rats. ^a*P* < 0.001 when negative group was compared with normal group; ^x*P* < 0.001, ^b*P* < 0.05 when test groups were compared with negative group

Histopathological observation

Histopathological observation of liver sections of the different group supports the hepatoprotective activity of the rind extract. The control group animal exhibited normal hepatic cell with defined cytoplasm, prominent nucleus, and normal hepatic vein [Figure 9a]. The CCl₄ intoxicated group showed lost of total hepatic architecture with necrosis, inflammation, crowding central vein in many areas [Figure 9b]. The others group pretreatment with Silymarin and rind extract (200, 400 mg/kg) showed the liver protection by minimizing inflammation, less necrosis and reducing crowding of central vein in a dose-dependent manner [Figure 9c-e].

DISCUSSION

Phytochemical are usually recognized as antioxidant which contain bioactive polyphenols, such as phenolic acid and flavonoids. Phenolic acid contains the major phenol containing groups which are served as a primary source of antioxidant or free radical scavenger, while polyphenols act as an antioxidant terminator by scavenging free radicals to form stable one.^[28,29] In current findings, it was confirmed that phenol and flavonoids are present in *G. dulcis* rind extract, which are responsible for scavenging free radicals in various antioxidant models.




Figure 9: Effect of methanolic fruit rind extract of *Garcinia dulcis* on liver. Liver section, 0.5 μ size, H and E, $\times 40$, (a) control group, exhibiting the normal liver architecture with normal hepatic cells, with define cytoplasm, prominent nucleus and normal central vein; (b) negative group, exhibiting total loss of hepatic architecture with necrosis, crowding of central vein; (c) standard group, exhibiting normal hepatic architecture with minimal necrosis, normal central vein, (d) low-dose test group (200 mg/kg) showing a pattern of reduced necrosis, (e) high-dose group, (400 mg/kg), is showing normal architecture with normal hepatic and normal central vein

DPPH evolves free radicals whenever it puts in solvents. The DPPH system is widely accepted tools to determine the free radical scavenging activities of antioxidants; it accepts an electron or hydrogen radical as stable molecules.^[30] The deductive capability of free radicals in DPPH system can be determined by decreasing absorbance at 517 nm, experimentally. The present findings revealed that *G. dulcis* rind extract has the potential to scavenge the DPPH free radicals in a dose-dependent manner.

The ABTS is relatively a recent free radical scavenging method which produces more drastic radicals, often used for screening complex antioxidant from the mixture such as plant extract, biological beverage, and fluids.^[31] In this method, the antioxidant potentiality is measured by loss of blue-green color of ABTS•+ when the antioxidant is putted into it. The antioxidant reduces ABTS•+ to stable ABTS and decolorize.^[32] Our finding regarding on removal of ABTS•+ free radicals is dose-dependent. Nitride oxide produces continuously inside the human body, intermediating the regulation of various metabolic processes. Excess production of NO• may cause several diseases such as inflammation, cancer, etc. NO• is reactive under aerobic conditions and usually react with oxygen to produce stable product, i.e., nitrate and nitrite by producing intermediates such as NO₂, N₂O₄, and N₃O₄.^[33] These free radicals cause alternating the functional behavior of cellular component. The free radical scavenging principle in NO• models is that the antioxidants present in extract compete with oxygen to react with nitride oxide and inhibits the production of intermediate compounds, Hence, protect the

cellular components.^[31] Nitride oxide can be generated by incubation of sodium nitroprusside in PBS at 25°C. The present investigation on NO• scavenging potentiality of *G. dulcis* rind extract was found to be dose dependent.

Our results showed that the methanolic extract of the *G. dulcis* possess significant antioxidant activity. The involvement of free radicals in the damage of liver has been investigated for many years. The free radical scavenging potentiality of the extract was also accompanied by the hepatic protection induced by CCl₄. In, liver CCl₄ is metabolically activated by cytochrome P450 mixed oxidase in endoplasmic reticulum to form CCl₃• free radicals which bind covalently to unsaturated lipid membrane, elevation of bilirubin level, reduction of total protein of serum and followed by liver enzyme marker level such as SGOT, SGPT, SALP, and finally hepatocyte damage.^[34,35]

The assessment of liver protection can be evaluated by determining the serum enzyme. Hepatocellular necrosis or cell membrane injury leads to elevate of serum SGOT and SGPT, released from liver to blood circulation. The assessment of SGPT is considered as a high index because it is found mainly in the liver and catalase conversion of alanine to pyruvate and glutamate, and release as similar manner to the circulation. Hence, it covered the 90% of total enzyme.^[36] Slight elevation of this enzyme is considered as myocardial infarction, jaundice, etc., SALP is related to functioning hepatocyte and elevation in the serum is indicated the biliary pressure.^[37] Our results show increased the level of serum enzymes in the negative group than the normal group. Administration of the methanolic fruit rind extract of *G. dulcis* and silymarin significantly reverses the elevated level of these enzymes, in a dose-dependent manner. Silymarin is well-known hepatoprotective agent derived from *Silybum marianum*. Hence, silymarin was used as the standard drug.^[38] The result indicated that the methanolic fruit rind extract of *G. dulcis* can ameliorate hepatic function of liver damage by CCl₄.

Alternation of total protein has been considered as one of the major factor associated with the hepatic dysfunction.^[39] The present study showed that the protein level was decreased in CCl₄ induced group compared to control group, which restored almost the normal range pretreatment with the *G. dulcis* extract a dose-dependent manner.

Bilirubin usually removed from the blood by liver through conjugation and secreted in the bile. It breakdown of the red blood cells and elevation of this is indicating the hepatic dysfunction.^[40] The current study found that the bilirubin concentration was elevated in CCl₄ induced group, compared to the control group and pretreatment with the extract restored the level of bilirubin in a dose-dependent manner.

CONCLUSION

The present finding revealed that the fruit rind extract of *G. dulcis* contains both phenolic and flavonoids compounds, possesses potent antioxidant activity *in vitro* models and a significant hepatoprotective activity. These findings indicate that the fruit can be used as functional fruit with any toxicity, and their uses should be considered as an adjunct therapeutic strategy to combat different hepatic disorder and other diseases caused by free radicals. In addition, further studies to separate and characterized the active compounds of *G. dulcis* and their mechanism toward the hepatoprotective are in progress.

Acknowledgments

The authors express sincere gratitude to Department of Pharmaceutical Science, Dibrugarh University for providing the Animal house facility. The authors also thank Prof. A. K. Rai and Banasri Mech, Research Scholar, Department of Life Sciences, Dibrugarh University for help and providing Microtome facility.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M. Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 2006;160:1-40.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 2007;39:44-84.
- Rajkumar V, Guha G, Kumar RA. Antioxidant and anti-neoplastic activities of *Picrorhiza kurroa* extracts. *Food Chem Toxicol* 2011;49:363-9.
- Wichi HP. Enhanced tumor development by butylated hydroxyanisole (BHA) from the prospective of effect on forestomach and oesophageal squamous epithelium. *Food Chem Toxicol* 1988;26:717-23.
- Sharma A, Chakraborti KK, Handa SS. Antihepatotoxic activity of some Indian herbal formulations as compared to silymarin. *Fitoterapia* 1991;62:229-35.
- Hiraganahalli BD, Chinampudur VC, Dethle S, Mundkinajeddu D, Pandre MK, Balachandran J, *et al.* Hepatoprotective and antioxidant activity of standardized herbal extracts. *Pharmacogn Mag* 2012;8:116-23.
- Kanter M, Coskun O, Budancamanak M. Hepatoprotective effects of *Nigella sativa* L and *Urtica dioica* L on lipid peroxidation, antioxidant enzyme systems and liver enzymes in carbon tetrachloride-treated rats. *World J Gastroenterol* 2005;11:6684-8.
- Sharma S, Sahu AN. Development, characterization, and evaluation of hepatoprotective effect of *Abutilon indicum* and *Piper longum* Phytosomes. *Pharmacognosy Res* 2016;8:29-36.
- Wiseman H, Halliwell B. Damage to DNA by reactive oxygen and nitrogen species: Role in inflammatory disease and progression to cancer. *Biochem J* 1996;313(Pt 1):17-29.
- Rao GM, Rao CV, Pushpangadan P, Shirwaikar A. Hepatoprotective effects of rubiadin, a major constituent of *Rubia cordifolia* Linn. *J Ethnopharmacol* 2006;103:484-90.
- Cordero Pérez P, Torres-González L, Aguirre-Garza M, Camara-Lemarrroy C, Guzmán-de la Garza F, Alarcón-Galván G, *et al.* Hepatoprotective effect of commercial herbal extracts on carbon tetrachloride-induced liver damage in Wistar rats. *Pharmacogn Res* 2013;5:150.
- Burkill IH. *A Dictionary of Economic Product of the Malay Peninsula*. 1st ed. London: Government of Straits Settlement; 1935.
- Gogoi N, Gogoi A, Neog B. Free radical scavenging activities of *Garcinia xanthochymus* Hook. f. and *Garcinia lanceifolia* Roxb. using various *in vitro* assay models. *Asian J Pharm Clin Res* 2015;9:138-41.
- Begum A, Barthakur SK, Sarma J. *Garcinia dulcis* (Roxburgh) Kurz [*Clusiaceae*]: A new distributional record for Assam, India. *Pleione* 2013;7:545-8.
- Kumaran A, Karunakaran RJ. *In vitro* antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *LWT Food Sci Technol* 2007;40:344-52.
- Hsu CY. Antioxidant activity of extract from *Polygonum aviculare* L. *Biol Res* 2006;39:281-8.
- Shirwaikar A, Shirwaikar A, Rajendran K, Punitha IS. *In vitro* antioxidant studies on the benzyl tetra isoquinoline alkaloid berberine. *Biol Pharm Bull* 2006;29:1906-10.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 1999;26:1231-7.
- Marcocci L, Maguire JJ, Droy-Lefaix MT, Packer L. The nitric oxide-scavenging properties of *Ginkgo biloba* extract EGB 761. *Biochem Biophys Res Commun* 1994;201:748-55.
- Srinivasan R, Chandrasekar MJ, Nanjan MJ, Suresh B. Antioxidant activity of *Caesalpinia digyna* root. *J Ethnopharmacol* 2007;113:284-91.
- OECD 2001-gudeline on acute oral toxicity (AOT). Environmental health and safety monograph series on testing and adjustment. No. 425.
- Singh K, Khanna AK, Chander R. Hepatoprotective activity of ellagic acid against carbon tetrachloride induced hepatotoxicity in rats. *Indian J Exp Biol* 1999;37:1025-6.
- Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am J Clin Pathol* 1957;28:56-63.
- King EJ, Kind RP. Alkaline phosphatase activity assay. *J Clin Pathol* 1954;7:332.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951;193:265-75.
- Malloy HT, Evelyn KA. The determination of bilirubin with the photoelectric colorimeter. *J Biol Chem* 1937;119:481-90.
- Galigher AE, Kozloff EN. *Essentials of Practical Microtechniques*. 2nd ed. Philadelphia: Lea and Febiger; 1971.
- Grimmer HR, Parbhoo V, McGrath RM. Antimutagenicity of polyphenol-rich fractions from sorghum bicolor grain. *J Sci Food Agric* 1992;59:251-6.
- Almulaiky YQ, Alshawafi WM, Al-Talhi HA, Zeyadi M, Anwar F, Al-abbasi FA, *et al.* Evaluation of the antioxidant potential and antioxidant enzymes of some Yemeni grape cultivars. *Free Radic Antioxid* 2017;7:74.
- Kalaivani T, Mathew L. Free radical scavenging activity from leaves of *Acacia nilotica* (L.) Wild. ex Delile, an Indian medicinal tree. *Food Chem Toxicol* 2010;48:298-305.
- Kumar RS, Raj Kapoor B, Perumal P. Antioxidant activities of *Indigofera cassioides* Rottl. Ex. DC. using various *in vitro* assay models. *Asian Pac J Trop Biomed* 2012;2:256-61.
- Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J* 2013;21:143-52.
- Jayakumari S, Arthanareswaran A, Vijayalakshmi A, Velraj M, Ravichandran V. Free radical scavenging activity of *Pisonia grandis* R. Br leaves. *Int J Pharm Educ Res* 2012;46:37.
- Kaplowitz N, Aw TY, Simon FR, Stolz A. Drug-induced hepatotoxicity. *Ann Intern Med* 1986;104:826-39.
- Kadiiska MB, Gladen BC, Baird DD, Dikalova AE, Sohal RS, Hatch GE, *et al.* Biomarkers of oxidative stress study: Are plasma antioxidants markers of CCl(4) poisoning? *Free Radic Biol Med* 2000;28:838-45.
- Achliya GS, Kotagale NR, Wadodkar SG, Dorle AK. Hepatoprotective activity of panchagavya ghrita against carbon tetrachloride induced hepatotoxicity in rats. *Indian J Pharmacol* 2003;35:308-11.
- Moss DW, Butterworth PJ. *Enzymology and Medicine*. London: Pitman Medical; 1974.
- Jose MA, Abraham A, Narmadha MP. Effect of silymarin in diabetes mellitus patients with liver diseases. *J Pharmacol Pharmacother* 2011;2:287-9.
- Charlton MR. Protein metabolism and liver disease. *Baillieres Clin Endocrinol Metab* 1996;10:617-35.
- Naik SR, Panda VS. Antioxidant and hepatoprotective effects of *Ginkgo biloba* phytosomes in carbon tetrachloride-induced liver injury in rodents. *Liver Int* 2007;27:393-9.